

PA 1236214

REC'D 11 NOV 2004

WIPO

PCT

GB 04/4278

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

October 15, 2004


THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/509,936

FILING DATE: October 10, 2003

**PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)**


**By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS**



W. Montgomery
W. MONTGOMERY
Certifying Officer

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: James T. FULLER
Title: NUCLEIC ACID CONSTRUCTS
Appl. No.: Unknown
Filing Date: 10/10/2003
Examiner: Unknown
Art Unit: Unknown



PROVISIONAL PATENT APPLICATION
TRANSMITTAL

Mail Stop PROVISIONAL PATENT APPLICATION
Commissioner for Patents
PO Box 1450
Alexandria, Virginia 22313-1450

Sir:

Transmitted herewith for filing under 37 C.F.R. § 1.53(c) is the provisional patent application of:

James T. FULLER
585 Science Drive
Madison, Wisconsin 53711

☐ Applicant claims small entity status under 37 CFR 1.27(c)(1).

Enclosed are:

- ☒ Specification and Abstract (117 pages).
- ☒ Informal drawings (13 sheets, Figures 1-13).
- ☒ Application Data Sheet (37 CFR 1.76).

The filing fee is calculated below:

	Rate	Fee Totals
Basic Fee	\$160.00	\$160.00
[] Small Entity Fees Apply (subtract ½ of above):	=	\$0.00
TOTAL FILING FEE:	=	\$160.00

- [X] A check in the amount of \$160.00 to cover the filing fee is enclosed.
- [] The required filing fees are not enclosed but will be submitted in response to the Notice to File Missing Parts of Application.
- [X] Further, Applicant, by and through his attorney of record, hereby expressly abandons the application as of the filing date of this letter. This is an abandonment of the application only, and is not to be construed as an abandonment of the invention disclosed in the application. It is respectfully requested that the Office acknowledge abandonment of the application as of the filing date of this letter in a communication mailed to the undersigned.
- [X] The Commissioner is hereby authorized to charge any additional fees which may be required regarding this application under 37 C.F.R. §§ 1.16-1.17, or credit any overpayment, to Deposit Account No. 19-0741. Should no proper payment be enclosed herewith, as by a check being in the wrong amount, unsigned, post-dated, otherwise improper or informal or even entirely missing, the Commissioner is authorized to charge the unpaid amount to Deposit Account No. 19-0741.

Please direct all correspondence to the undersigned attorney or agent at the address indicated below.

Respectfully submitted,

Date Oct-10, 2003

FOLEY & LARDNER
Customer Number: 27476
Telephone: (202) 672-5483
Facsimile: (202) 672-5399

By Richard C. Peet (28,665)
for Richard C. Peet
Attorney for Applicant
Registration No. 35,792

Application Data Sheet

Application Information

Application Type::	Provisional
Subject Matter::	Utility
Suggested classification::	
Suggested Group Art Unit::	
CD-ROM or CD-R?::	None
Computer Readable Form (CRF)?::	No
Title::	NUCLEIC ACID CONSTRUCTS
Attorney Docket Number::	036481-0141
Request for Early Publication?::	No
Request for Non-Publication?::	No
Suggested Drawing Figure::	1
Total Drawing Sheets::	13
Small Entity?::	Yes
Petition included?::	No
Secrecy Order in Parent Appl.?::	No

Applicant Information

Applicant Authority Type::	Inventor
Primary Citizenship Country::	US
Status::	Full Capacity
Given Name::	James T.
Family Name::	FULLER
City of Residence::	Madison
Country of Residence::	Wisconsin
Street of mailing address::	585 Science Drive
City of mailing address::	Madison

State or Province of mailing **WI**
address::
Postal or Zip Code of mailing **53711**
address::

Correspondence Information

Correspondence Customer Number:: **27476**
E-Mail address:: **PTOMailWashington@FoleyLaw.com**

Representative Information

Representative Customer Number::	22428	
---	--------------	--

Domestic Priority Information

Application::	Continuity Type::	Parent Application::	Parent Filing Date::

Foreign Priority Information

Country::	Application number::	Filing Date::	Priority Claimed::

Assignee Information

Assignee name::

Atty. Dkt. No.: 036481/0141

U.S. PATENT PROVISIONAL APPLICATION

for

NUCLEIC ACID CONSTRUCTS

Inventors: James T. FULLER

NUCLEIC ACID CONSTRUCTS

FIELD OF THE INVENTION

The invention relates to the fields of molecular biology and immunology and generally to reagents useful in nucleic acid immunisation techniques. More specifically, the invention relates to nucleic acid constructs for the expression of antigenic polypeptides, and to nucleic acid immunisation strategies using such reagents.

BACKGROUND OF THE INVENTION

Gene therapy and nucleic acid immunisation are promising approaches for the treatment and prevention of both acquired and inherited diseases. These techniques provide for the transfer of a desired nucleic acid into a subject with subsequent *in vivo* expression. Transfer can be accomplished by transfecting the subject's cells or tissues *ex vivo* and reintroducing the transformed material into the host. Alternatively, the nucleic acid can be administered *in vivo* directly to the recipient.

Each of these techniques requires efficient expression of the nucleic acid in the transfected cell, to provide a sufficient amount of the therapeutic or antigenic gene product. Several factors are known to affect the levels of expression obtained, including transfection efficiency, and the efficiency with which the gene or sequence of interest is transcribed and the mRNA translated.

A number of expression systems have been described in the art, each of which typically consists of a vector containing a gene or nucleotide sequence of interest operably linked to expression control sequences. These control sequences include transcriptional promoter sequences and transcriptional start and termination sequences.

Commonly used promoters for mammalian cell expression systems include the SV40 early promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter (Chapman *et al* (1991) *Nucl. Acids Res.* 19:3979-3986), the mouse mammary tumour virus long terminal repeat (LTR) promoter, the adenovirus major late promoter (Ad MLP) and the herpes simplex virus (HSV) promoter, among others. Nonviral promoters, such as a promoter derived from the murine metallothionein gene are also commonly used.

Expression systems often include transcriptional modulator elements, referred to as "enhancers". Enhancers are broadly defined as a cis-acting agent, which when operably linked to a promoter/gene sequence, will increase transcription of that gene sequence. Enhancers can function from positions that are much further away from a sequence of interest than other expression control elements (e.g. promoters), and can operate when positioned in either orientation relative to the sequence of interest (Banerji *et al.* (1981) *Cell* 27:299-308, deVilleirs *et al.* (1981) *Nucl. Acids Res* 9: 6251-6264). Enhancers have been identified from a number of viral sources, including polyoma virus, BK virus, cytomegalovirus (CMV), adenovirus, simian virus 40 (SV40), Moloney sarcoma virus, bovine papilloma virus and Rous sarcoma virus (deVilleirs *et al supra*, Rosenthal *et al.* (1983) *Science* 222:749-755, Hearing *et al.* (1983) *Cell* 33:695-703, Weeks *et al.* (1983) *Mol. Cell. Biol.* 3:1222-1234, Levinson *et al.* (1982) *Nature* 295: 568-572, and Luciw *et al.* (1983) *Cell* 33: 705-716).

A number of expression systems for nucleic acid immunisation and gene therapy make use of the hCMV immediate early promoter. See eg US Patent Nos. 5168062 and 5385839 to Stinski, and EP Patent Specification 0323997 B1. Expression vectors using the hCMV

immediate early promoter include for example, pWRG7128 (Roy *et al*, Vaccine 19, 764-778, 2001), and pBC12/CMV and pJW4303 which are mentioned in WO 95/20660. Chapman *et al* (1991) report reduced levels of expression from the hCMV immediate early promoter in the absence of hCMV Intron A.

SUMMARY OF THE INVENTION

A nucleic acid construct has been developed using manipulated viral promoter/expression sequences, that provides enhanced expression of heterologous coding sequences in host cells. The construct is suitable for efficient expression of antigen-encoding genes, and can therefore be used in nucleic acid immunisation. In particular, the construct can be provided on carrier particles, for use in particle-mediated nucleic acid immunisation.

Accordingly, the present invention provides a nucleic acid construct comprising a chimeric promoter sequence and a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter, wherein the chimeric promoter sequence comprises:

- (a) a hCMV immediate early promoter sequence;
- (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
- (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

The invention also provides:

- a nucleic acid construct comprising:

(i) a chimeric promoter sequence which comprises:

(a) a hCMV immediate early promoter sequence;

(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene; and

(ii) a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter; and

(iii) (a) a non-translated leader sequence which is derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2gD antigen sequence and which is in operable linkage with the chimeric promoter; and/or

(b) an enhancer sequence which is derived from a 3' untranslated region (UTR) of a HBsAg sequence, or a 3'UTR of a simian CMV immediate early gene sequence and which is in operable linkage with the chimeric promoter, wherein the enhancer sequence is downstream of the cloning site;

- a nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen sequence; and

(iii) a coding sequence operably linked to (i) and (ii)

wherein the coding sequence is heterologous to the non-translated leader sequence;

- a nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a coding sequence operably linked to the promoter sequence (i); and

(iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence, and the coding sequence (ii) is heterologous to the 3' enhancer sequence;

- a method of obtaining expression in mammalian cells of a polypeptide of interest, which method comprises transferring into said cells a nucleic acid construct of the invention, the construct including a coding sequence encoding the polypeptide;

- coated particles, suitable for delivery from a particle-mediated delivery device, which particles comprise carrier particles coated with a nucleic acid construct of the invention, the construct including a coding sequence encoding the polypeptide;

- a dosage receptacle for a particle mediated delivery device comprising the coated particles;

- a particle mediated delivery device loaded with the coated particles;

- a method of nucleic acid immunisation comprising administering to a subject an effective amount of the coated particles in which particles the coding sequence encodes an antigen;

- a purified isolated chimeric promoter sequence which comprises:

(a) a hCMV immediate early promoter sequence;

(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

These and other objects, aspects, embodiments and advantages of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

Accordingly, the present invention provides a nucleic acid construct comprising a chimeric promoter sequence and a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter, wherein the chimeric promoter sequence comprises:

- (a) a hCMV immediate early promoter sequence;
- (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
- (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

In one embodiment, the hCMV immediate early promoter sequence (a) is SEQ ID No.1.

In another embodiment, the exon sequence (b) is SEQ ID No.2.

In another embodiment, the heterologous intron (c) comprises a sequence selected from the group consisting of rat insulin gene intron A sequence, chicken keratin gene intron A sequence and chicken cardiac actin gene intron A sequence. In another embodiment, the rat insulin gene intron A sequence comprises SEQ ID No. 3. In another embodiment, the chimeric promoter sequence is SEQ ID No. 4.

In another embodiment, the nucleic acid construct comprises a polyadenylation sequence. In a further embodiment, the polyadenylation sequence is derived from a polyadenylation sequence of a gene selected from the group consisting of rabbit β -globin gene, human papilloma virus (HPV) early or late genes, the HSV-2gB gene, a simian CMV immediate early gene and HSVgD late gene. In a preferred embodiment, the

polyadenylation sequence is selected from the group consisting of SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12 and SEQ ID NO:13.

This particular nucleic acid construct may further comprise a signal peptide. In one embodiment, the signal peptide is selected from the group consisting of human tissue plasminogen activator signal peptide (hTPAsp), aprotinin signal peptide, tobacco extensin signal peptide and chicken lysozyme signal peptide.

In another embodiment, a coding sequence is provided in the cloning site. In a preferred embodiment the coding sequence encodes an antigen. In one embodiment, the antigen is an antigen of a viral, bacterial, parasitic or fungal pathogen. In a preferred embodiment, the antigen is HBsAg.

This particular construct may be a plasmid vector. Furthermore, the nucleic acid in this construct is DNA.

In another aspect of the present invention, another nucleic acid construct is provided, which comprises:

(i) a chimeric promoter sequence which comprises:

(a) a hCMV immediate early promoter sequence;

(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene; and

(ii) a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter; and

(iii) (a) a non-translated leader sequence which is derived from HBVpreS2 antigen sequence, HBV e-antigen sequence or HSV type 2gD antigen sequence and which is in operable linkage with the chimeric promoter; and/or

(b) an enhancer sequence which is derived from a 3' untranslated region (UTR) of a HBsAg sequence, or a 3'UTR of a simian CMV immediate early gene sequence and which is in operable linkage with the chimeric promoter, wherein the enhancer sequence is downstream of the cloning site.

In one embodiment, the hCMV immediate early promoter sequence (a) is SEQ ID No.1.

In another embodiment, the exon sequence (b) is SEQ ID No.2.

In another embodiment, the heterologous intron (c) comprises a sequence selected from the group consisting of rat insulin gene intron A sequence, chicken keratin gene intron A sequence and chicken cardiac actin gene intron A sequence. In a preferred embodiment, the rat insulin gene intron A sequence comprises SEQ ID No. 3.

In yet another embodiment, the chimeric promoter sequence is SEQ ID No. 4.

Also, this construct may comprise a polyadenylation sequence. In one embodiment, the polyadenylation sequence is derived from a polyadenylation sequence of a gene selected from the group consisting of a rabbit β -globin gene, human papilloma virus (HPV) early or late genes, the HSV-2gB gene, a simian CMV immediate early gene and HSVgD late gene. In a preferred embodiment, the polyadenylation

sequence is selected from the group consisting of SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12 and SEQ ID NO:13.

This construct may further comprise a signal peptide. In one embodiment, the signal peptide is selected from the group consisting of human tissue plasminogen activator signal peptide (hTPAsp), aprotinin signal peptide, tobacco extensin signal peptide and chicken lysozyme signal peptide.

In another embodiment, the construct comprises a coding sequence in the cloning site.

In another embodiment, a coding sequence is provided in the cloning site. In a preferred embodiment the coding sequence encodes an antigen. In one embodiment, the antigen is an antigen of a viral, bacterial, parasitic or fungal pathogen. In a preferred embodiment, the antigen is HBsAg.

This particular construct may be a plasmid vector. Furthermore, the nucleic acid in this construct is DNA.

In yet another embodiment, this nucleic acid construct has the sequence in SEQ ID No. 14.

In a further embodiment, the non-translated leader sequence in the nucleic acid construct is selected from the group consisting of SEQ ID No. 5, SEQ ID No. 6 and SEQ ID NO:7.

In another embodiment, the enhancer sequence in the nucleic acid construct is selected from the group consisting of SEQ ID No. 8 and SEQ ID No. 9.

In another aspect of the present invention, a method of obtaining expression in mammalian cells of a polypeptide of interest is provided. This method comprises transferring into said cells a nucleic acid construct comprising a chimeric promoter sequence and a coding sequence encoding the polypeptide in operable linkage with the chimeric promoter, wherein the chimeric promoter sequence comprises:

(a) a hCMV immediate early promoter sequence;

(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

In one embodiment, the construct is delivered directly into a subject. In another embodiment, the construct is delivered by injection, transdermal particle delivery, inhalation, topically, orally, intranasally or transmucosally. In a preferred embodiment, the construct is delivered by needleless injection.

In yet another embodiment, the construct is delivered *ex vivo* into cells taken from a subject, and the cells are reintroduced into the subject.

In another embodiment, the nucleic acid construct is coated onto carrier particles.

The present invention also encompasses coated particles, suitable for delivery from a particle-mediated delivery device, which particles comprise carrier particles coated with a nucleic acid construct wherein the construct comprises a chimeric promoter sequence and a coding

sequence in operable linkage with the chimeric promoter, and wherein the chimeric promoter sequence comprises:

- (a) a hCMV immediate early promoter sequence;
- (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
- (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

In one embodiment, the carrier particles are gold or tungsten.

In yet another aspect of the invention is a dosage receptacle for a particle mediated delivery device comprising the coated particles described herein. In one embodiment, a particle mediated delivery device is loaded with such coated particles. In a preferred embodiment, the particle mediated delivery device is a needleless syringe.

The present invention also provides a method of nucleic acid immunization, comprising administering to a subject an effective amount of coated particles, which particles are suitable for delivery from a particle-mediated delivery device, the particles comprising carrier particles coated with a nucleic acid construct, wherein the construct comprises a chimeric promoter sequence and a coding sequence encoding an antigen in operable linkage with the chimeric promoter, and wherein the chimeric promoter sequence comprises

- (a) a hCMV immediate early promoter sequence;
- (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

Also provided in the present invention is a nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen sequence; and

(iii) a coding sequence operably linked to (i) and (ii)

wherein the coding sequence is heterologous to the non-translated leader sequence. In one embodiment, the promoter sequence (i) is selected from the group consisting of hCMV immediate early promoter sequence, Pseudorabies virus promoter sequence and Rous sarcoma virus promoter sequence.

In another embodiment, the coding sequence encodes an antigen. In another embodiment, the antigen is an antigen of a viral, bacterial, parasitic or fungal pathogen.

In another embodiment, the nucleic acid construct is a plasmid vector.

In yet another aspect of the present, a method of obtaining expression in mammalian cells of a polypeptide of interest is provided, which method comprises transferring into said cells a nucleic acid construct comprising: (i) a promoter sequence; (ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen sequence; and (iii) a coding

sequence encoding the polypeptide operably linked to (i) and (ii), wherein the coding sequence is heterologous to the non-translated leader sequence.

In one embodiment, the construct is delivered directly into a subject. In another embodiment, the construct is delivered by injection, transdermal particle delivery, inhalation, topically, orally, intranasally or transmucosally. In a preferred embodiment, the construct is delivered by needleless injection.

In yet another embodiment, the construct is delivered *ex vivo* into cells taken from a subject, and the cells are reintroduced into the subject.

In another embodiment, the nucleic acid construct is coated onto carrier particles.

The present invention also encompasses coated particles, suitable for delivery from a particle-mediated delivery device, which particles comprise carrier particles coated with a nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen sequence; and

(iii) a coding sequence encoding a polypeptide of interest operably linked to (i) and (ii)

wherein the coding sequence is heterologous to the non-translated leader sequence.

In one embodiment, these carrier particles are gold or tungsten.

In yet another aspect of the invention is a dosage receptacle for a particle mediated delivery device comprising the coated particles described herein. In one embodiment, a particle mediated delivery device is loaded with such coated particles. In a preferred embodiment, the particle mediated delivery device is a needleless syringe.

In one other aspect of the present invention is a method of nucleic acid immunization comprising administering to a subject an effective amount of coated particles which are suitable for delivery from a particle mediated delivery device, the particles comprising carrier particles coated with a nucleic acid construct comprising

- (i) a promoter sequence;
- (ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2gD antigen sequence; and
- (iii) a coding sequence encoding an antigen operably linked to (i) and (ii) wherein the coding sequence is heterologous to the non translated leader sequence.

In yet another aspect of the present invention is provided a nucleic acid construct comprising:

- (i) a promoter sequence;
- (ii) a coding sequence operably linked to the promoter sequence (i); and

(iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence, and the coding sequence (ii) is heterologous to the 3'enhancer sequence. In one embodiment of this construct, the promoter sequence (i) is selected from the group consisting of hCMV immediate early promoter sequence, Pseudorabies virus promoter sequence and Rous sarcoma virus promoter sequence.

In another embodiment, the coding sequence encodes an antigen. In yet another embodiment, the antigen is an antigen of a viral, bacterial, parasitic or fungal pathogen.

In another embodiment, the nucleic acid construct is a plasmid vector.

The present invention also provides a method of obtaining expression in mammalian cells of a polypeptide of interest, which method comprises transferring into said cells a nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a coding sequence encoding the polypeptide operably linked to the promoter sequence (i); and

(iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene

sequence and the coding sequence (ii) is heterologous to the 3'enhancer sequence.

In one embodiment, the construct is delivered directly into a subject. In one embodiment, the construct is delivered by injection, transdermal particle delivery, inhalation, topically, orally, intranasally or transmucosally. In a preferred embodiment, the construct is delivered by needleless injection.

In yet another embodiment, the construct is delivered *ex vivo* into cells taken from a subject, and the cells are reintroduced into the subject.

In another embodiment, the nucleic acid construct is coated onto carrier particles.

The present invention also encompasses coated particles, suitable for delivery from a particle-mediated delivery device, which particles comprise carrier particles coated with a nucleic acid construct comprising:

- (i) a promoter sequence;
- (ii) a coding sequence encoding a polypeptide of interest operably linked to the promoter sequence (i); and
- (iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence and the coding sequence (ii) is heterologous to the 3' enhancer sequence.

In one embodiment, the carrier particles are gold or tungsten.

In yet another embodiment, the dosage receptacle for a particle mediated delivery device comprising coated particles is provided. Also provided in the present invention is a particle mediated delivery device loaded with such coated particles. In a preferred embodiment, the particle mediated delivery device is a needleless syringe.

The present invention also provides a method of nucleic acid immunisation comprising administering to a subject an effective amount of coated particles which particles are suitable for delivery from a particle mediated delivery device, the particles comprising carrier particles coated with a nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a coding sequence encoding an antigen operably linked to the promoter sequence (c); and

(iii) an enhancer sequence 3' of and operably linked to the coding sequence (iii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence and the coding sequence (ii) is heterologous to the 3'enhancer sequence.

In yet another aspect of the present invention is a purified isolated chimeric promoter sequence which comprises:

(a) a hCMV immediate early promoter sequence;

(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the levels of expression of hepatitis B virus surface antigen (HBsAg) obtained using various plasmid expression vectors.

Figure 2 shows the effect of intron inclusion on expression of HBsAg in SSC15 cells and of beta-gal in B16 cells (average of three expts).

Figure 3 shows the effect of rat insulin intron A and HBV3'UTR on expression of beta-gal in SSC15 cells (average of three expts).

Figure 4 shows the effect of rat insulin intron A and HBV3'UTR on expression of HSVgD in SSC15 cells (average of three expts).

Figure 5 shows the effect of rat insulin intron A and HBV3'UTR on expression of SEAP in SSC15 and B16 cells (three repetitions per cell line).

Figure 6 shows the ability of heterologous signal peptides to direct secretion of SEAP or hFc fragment in B16 cells.

Figure 7 illustrates levels of antibodies detected in the sera of mice immunised with antigen-encoding nucleic acids contained in a variety of plasmid expression vectors.

Figure 8 is a diagrammatic representation of pJV expression vector.

Figure 9 is a diagrammatic representation of pJV7389.

Figure 10 is a diagrammatic representation of pJV7400.

Figure 11 is a diagrammatic representation of pJV7468.

Figure 12 is a diagrammatic representation of pJV7563.

Figure 13 sets out the base composition for pJV7563.

DETAILED DESCRIPTION

Before describing the present invention in detail, it is to be understood that this invention is not limited to particularly exemplified molecules or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting. In addition, the practice of the present invention will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology, recombinant DNA techniques and immunology all of which are within the ordinary skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., 1984); *A Practical Guide to Molecular Cloning* (1984); and *Fundamental Virology*, 2nd Edition, vol. I & II (B.N. Fields and D.M. Knipe, eds.).

All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

It must be noted that, as used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the content clearly dictates otherwise.

A. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although a number of methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred materials and methods are described herein.

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

The term "nucleic acid immunization" is used herein to refer to the introduction of a nucleic acid molecule encoding one or more selected antigens into a host cell for the *in vivo* expression of the antigen or antigens. The nucleic acid molecule can be introduced directly into the recipient subject, such as by standard intramuscular or intradermal injection; transdermal particle delivery; inhalation; topically, or by oral, intranasal or mucosal modes of administration. The molecule alternatively can be introduced *ex vivo* into cells which have been removed from a subject. In this latter case, cells containing the nucleic acid molecule of interest are re-introduced into the subject such that an immune response can be mounted against the antigen encoded by the

nucleic acid molecule. The nucleic acid molecules used in such immunization are generally referred to herein as "nucleic acid vaccines."

By "core carrier" is meant a carrier on which a guest nucleic acid (e.g., DNA, RNA) is coated in order to impart a defined particle size as well as a sufficiently high density to achieve the momentum required for cell membrane penetration, such that the guest molecule can be delivered using particle-mediated techniques (see, e.g., U.S. Patent No. 5,100,792). Core carriers typically include materials such as tungsten, gold, platinum, ferrite, polystyrene and latex. See e.g., *Particle Bombardment Technology for Gene Transfer*, (1994) Yang, N. ed., Oxford University Press, New York, NY pages 10-11.

By "needleless syringe" is meant an instrument which delivers a particulate composition transdermally without the aid of a conventional needle to pierce the skin. Needleless syringes for use with the present invention are discussed herein.

The term "transdermal" delivery intends intradermal (e.g., into the dermis or epidermis), transdermal (e.g., "percutaneous") and transmucosal administration, i.e., delivery by passage of an agent into or through skin or mucosal tissue. See, e.g., *Transdermal Drug Delivery: Developmental Issues and Research Initiatives*, Hadgraft and Guy (eds.), Marcel Dekker, Inc., (1989); *Controlled Drug Delivery: Fundamentals and Applications*, Robinson and Lee (eds.), Marcel Dekker Inc., (1987); and *Transdermal Delivery of Drugs*, Vols. 1-3, Kydonieus and Berner (eds.), CRC Press, (1987). Thus, the term encompasses delivery from a needleless syringe as described in U.S. Patent No. 5,630,796, as well as particle-mediated delivery as described in U.S. Patent No. 5,865,796.

A "polypeptide" is used in its broadest sense to refer to a compound of two or more subunit amino acids, amino acid analogs, or other peptidomimetics. The subunits may be linked by peptide bonds or by other bonds, for example ester, ether, etc. As used herein, the term "amino acid" refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is typically called a polypeptide or a protein.

An "antigen" refers to any agent, generally a macromolecule, which can elicit an immunological response in an individual. The term may be used to refer to an individual macromolecule or to a homogeneous or heterogeneous population of antigenic macromolecules. As used herein, "antigen" is generally used to refer to a protein molecule or portion thereof which contains one or more epitopes. For the purposes of the present invention, antigens can be obtained or derived from any appropriate source. Furthermore, for the purposes of the present invention, an "antigen" includes a protein having modifications, such as deletions, additions and substitutions (generally conservative in nature) to the native sequence, so long as the protein maintains sufficient immunogenicity. These modifications may be deliberate, for example through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the antigens.

An "immune response" against an antigen of interest is the development in an individual of a humoral and/or a cellular immune response to that antigen. For purposes of the present invention, a "humoral immune response" refers to an immune response mediated by

antibody molecules, while a "cellular immune response" is one mediated by T-lymphocytes and/or other white blood cells.

The terms "nucleic acid molecule" and "polynucleotide" are used interchangeably herein and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown. Non-limiting examples of polynucleotides include a gene, a gene fragment, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers.

A polynucleotide is typically composed of a specific sequence of four nucleotide bases: adenine (A); cytosine (C); guanine (G); and thymine (T) (uracil (U) for thymine (T) when the polynucleotide is RNA). Thus, the term nucleic acid sequence is the alphabetical representation of a polynucleotide molecule. This alphabetical representation can be input into databases in a computer having a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

A "vector" is capable of transferring nucleic acid sequences to target cells (e.g., viral vectors, non-viral vectors, particulate carriers, and liposomes). Typically, "vector construct," "expression vector," and "gene transfer vector," mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors. A "plasmid" is a vector in the form of an extrachromosomal genetic element.

A nucleic acid sequence which "encodes" a selected antigen is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. For the purposes of the invention, such nucleic acid sequences can include, but are not limited to, cDNA from viral, procaryotic or eucaryotic mRNA, genomic sequences from viral or procaryotic DNA or RNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence.

A "promoter" is a nucleotide sequence which initiates and regulates transcription of a polypeptide-encoding polynucleotide. Promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), repressible promoters (where expression of a polynucleotide sequence operably linked to the promoter is repressed by an analyte, cofactor, regulatory protein, etc.), and constitutive promoters. It is intended that the term "promoter" or "control element" includes full-length promoter regions and functional (*e.g.*, controls transcription or translation) segments of these regions.

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given promoter operably linked to a nucleic acid sequence is capable of effecting the expression of that sequence when the proper enzymes are present. The promoter need not be contiguous with the sequence, so long as it functions to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed

sequences can be present between the promoter sequence and the nucleic acid sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

"Recombinant" is used herein to describe a nucleic acid molecule (polynucleotide) of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation is not associated with all or a portion of the polynucleotide with which it is associated in nature and/or is linked to a polynucleotide other than that to which it is linked in nature. Two nucleic acid sequences which are contained within a single recombinant nucleic acid molecule are "heterologous" relative to each other when they are not normally associated with each other in nature.

Homologues of polynucleotides are referred to herein. Typically a polynucleotide which is homologous to another polynucleotide is at least 70% homologous to the polynucleotide, preferably at least 80 or 90% and more preferably at least 95%, 97% or 99% homologous thereto. Methods of measuring homology are well known in the art and it will be understood by those of skill in the art that in the present context, homology is calculated on the basis of nucleic acid identity. Such homology may exist over a region of at least 15, preferably at least 30, for instance at least 40, 60 or 100 or more contiguous nucleotides.

Methods of measuring polynucleotide homology or identity are known in the art. For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology (e.g. used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* 12, p387-395).

The PILEUP and BLAST algorithms can also be used to calculate homology or line up sequences (typically on their default settings), for

example as described in Altschul S.F. (1993) J Mol Evol 36:290-300;
Altschul, S, F *et al* (1990) J Mol Biol 215:403-10.

Software for performing BLAST analysis is publicly available through the National Centre for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al*, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89:10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5787. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by

chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

The homologues typically hybridize with the relevant polynucleotide at a level significantly above background. The signal level generated by the interaction between the homologue and the polynucleotide is typically at least 10 fold, preferably at least 100 fold, as intense as "background hybridisation". The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ^{32}P . Selective hybridisation is typically achieved using conditions of medium to high stringency, (for example, 0.03M sodium chloride and 0.003M sodium citrate at from about 50 EC to about 60 EC.

Stringent hybridization conditions can include 50% formamide, 5x Denhardt's Solution, 5x SSC, 0.1% SDS and 100 $\mu\text{g/ml}$ denatured salmon sperm DNA and the washing conditions can include 2x SSC, 0.1% SDS at 37EC followed by 1x SSC, 0.1% SDS at 68EC. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook *et al.*, *supra*.

The homologue may differ from a sequence in the relevant polynucleotide by less than 3, 5, 10, 15, 20 or more mutations (each of which may be a substitution, deletion or insertion). These mutations may be measured over a region of at least 30, for instance at least 40, 60 or 100 or more contiguous nucleotides of the homologue. Where a polynucleotide encodes a polypeptide, substitutions preferably create "conservative" changes in the amino acid encoded. These are defined according to the following table. Amino acids in the same block in the

second column and preferably in the same line in the third column may be substituted for each other in conservative changes.

ALIPHATIC	Non-Polar	G A P
		I L V
	Polar-uncharged	C S T M
		N Q
	Polar-charged	D E
		K R
AROMATIC		H F W Y

The terms "individual" and "subject" are used interchangeably herein to refer to any member of the subphylum cordata, including, without limitation, humans and other primates, including non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like. The terms do not denote a particular age. Thus, both adult and newborn individuals are intended to be covered. The methods described herein are intended for use in any of the above vertebrate species, since the immune systems of all of these vertebrates operate similarly.

B. General Overview

The invention is concerned with nucleic acid constructs which allow efficient expression of heterologous coding sequences, and in particular antigen-encoding genes, in host cells. More specifically, the

invention provides nucleic acid constructs comprising, or in some embodiments, consisting essentially of a chimeric promoter sequence and a cloning site, such that when a coding sequence is inserted in the cloning site, the coding sequence is in operable linkage with the chimeric promoter.

The chimeric promoter comprises, or in some embodiments consists essentially of :

- (a) a hCMV immediate early promoter sequence;
- (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
- (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

The hCMV immediate early promoter sequence (a) may comprise:

- (i) a native hCMV immediate early promoter sequence;
- (ii) a functional homologous variant thereof; or
- (iii) a functional fragment of (i) or (ii).

In general sequence (a) comprises about 100 to 600, preferably 200 to 600, for instance 400 to 600 nucleotides. Typically sequence (a) comprises the sequences present in (i) which bind transcription factors or the RNA polymerase, or instead of any of these sequences, homologues of these sequences able to bind the same transcription factors and RNA polymerase. Typically such sequences or their homologues are present in promoter sequence (a) in the same order and/or substantially the same relative spacing as in (i).

Generally, (i) comprises at least from nucleotides -100 to -1, typically -150 to -1, for example -500 to -1 or -600 to -1 of the hCMV major immediate early gene. Sequence (i) typically comprises the hCMV core promoter sequence and may also include one or more enhancer elements present in hCMV immediate early promoter. For example, (i) may comprise from nucleotides -118 to -1, or -524 to -1 as in US 6218140, or from nucleotides -62 to 1 or -465 to -1 as in US 538 5839.

Generally (i) includes a TATA box or CAAT box commonly found in promoter sequences. Preferably the sequence includes one or more of the repeat sequences in the hCMV immediate early promoter.

In a preferred embodiment, (i) comprises SEQ ID NO: 1.

A hCMV immediate early promoter sequence can be obtained using known methods. A native hCMV immediate early promoter can be isolated directly from a sample of the virus, using standard techniques. US 5385839, for example, describes the cloning of a hCMV promoter region. The sequence of a hCMV immediate early promoter is available at Genbank #M60321 (hCMV Towne strain) and X17403 (hCMV Ad169 strain). A native sequence could therefore be isolated by PCR using PCR primers based on the known sequence. See e.g. Sambrook *et al*, *supra*, for a description of techniques used to obtain and isolate DNA. A suitable hCMV promoter sequence could also be isolated from an existing plasmid vector. Promoter sequences can also be produced synthetically.

A functional variant (ii) or fragment (iii) is generally one which retains and/or complements the activity of the native promoter (i). Typically this activity is the ability to cause (including initiating and regulating) transcription of an operably linked polynucleotide, in particular the hCMV major immediate early gene. In one embodiment, the variant

or fragment would be able to complement the activity of the native promoter in a hCMV virus, for example allowing the virus to retain the ability to infect and/or replicate in cells.

A homologous variant (ii) or fragment (iii) can be assayed for the ability to retain and/or complement the activity of (i). For example, a variant or fragment may be assayed for ability to restore functionality (such as infection and/or replication ability) to mutant hCMV in which the native hCMV immediate early promoter is defective.

A homologous variant (ii) or fragment (iii) may be tested for utility using the Comparative Expression Assay below. The test promoter sequence is swapped into the base vector in place of the native hCMV immediate early promoter. Typically, a functional variant or fragment allows at least 50%, for instance, 60, 70, 80, 90% or more of the expression provided using the base vector. Generally expression is provided in at least one but preferably two reference cell types. Typically, the reference cells are mammalian HEK 293T, CHO, HeLa, BHK, 3T3 or COS cells.

Additionally or alternatively, promoter sequence may be tested in the Comparative Immunogenicity Assay below. Test promoter sequence is swapped into the base vector in place of the native hCMV immediate early promoter. A functional promoter sequence provides antibody titres that are at least as high as or higher than those achieved by the base vector with at least one, preferably two antigens. Preferably antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBsAg, HSV 2gD and flu-M2 antigens. According to this assay, a functional homologous variant (ii) or functional fragment (iii) of native promoter sequence (i) is one which allows the highest antibody titres achieved by the native sequence.

As mentioned above, the construct may comprise exon sequence (b) which comprises sequence derived from exon 1 and exon 2 of the hCMV major immediate early gene. Exons are coding sequences, which in nature are generally separated by introns. In the native hCMV major immediate early gene, exons 1 and 2 are usually separated by the native intron A. In the present chimeric construct exon 2 sequence is generally positioned 3' of exon 1 sequence, without intervening intron sequence, so that the exon 1 and exon 2 sequences are contiguous.

Exon sequence (b) may comprise:

- (i) native exon sequence, typically exon 1 and (whole or partial) exon 2;
- (ii) a homologous variant of (i) which is functional; or
- (iii) a functional fragment of (i) or (ii).

Sequence (i) may comprise from about 50 to 100% of the native hCMV major immediate early gene exon 1 sequence, for example, 60 to 90% or 70 to 80%. Typically at least 50% of the natural exon 1 sequence is present, such as 60%, 70%, 80%, 90% or more. Exon sequence (b) also comprises at least a part of exon 2 sequence. In sequence (i), typically 2 or more bases of native exon 2, for example 2 to 9, 2 to 7 or 3 to 5 bases are present. Up to and including 100% of natural exon 2 sequence, for example 5 to 95%, 20 to 80% or 40 to 60% of natural exon 2 sequence may be present. Typically the homologous variant has any of the above lengths mentioned for the native sequence.

Preferably (i) comprises SEQ ID No. 2.

Suitable exon sequence (b) can be obtained using known methods. See e.g. Sambrook *et al.*, *supra*, for a description of techniques used to obtain and isolate DNA. Native hCMV major immediate early gene sequence can be isolated directly from a sample of the virus, using standard techniques (see for example, MacLean, A (1998) "Preparation of HSV-DNA and Production of Infectious Virus" in Herpes Simplex Virus Protocols S. Brown, A Maclean, editors, Humana Press, Totowa, NJ, pp.19-26). The sequence of a hCMV major immediate early gene, including the location of exon 1 and exon 2, is available at Genbank #M60321, X17403. Native exon 1 and 2 sequences could therefore be isolated by cutting the native major gene sequence at appropriate restriction sites or by PCR using PCR primers based on the known sequence. Suitable exon sequences could alternatively be isolated from an existing plasmid vector, such as pWRG7128. Exon sequences can also be produced synthetically, rather than cloned. Variant sequences can readily be constructed by routine methodologies such as site-directed mutagenesis.

Generally the exon sequence will, when present in the construct of the invention, enhance expression, typically causing comparable enhancement to the native exon 1 and exon 2 sequence (i) mentioned above.

Exon sequence may be assayed for functionality using the Comparative Expression Assay below. Test exon sequence is swapped into the base vector in place of the exon sequence already present. Generally exon sequence is functional if the sequence does not abrogate expression but preferably increases expression in at least one but preferably two reference cell types when compared to the base vector. Typically the reference cells are mammalian HEK293T, CHO, HeLa, BHK,

3T3 or COS cells. Preferably expression increases by at least 5%, 10%, 20%, 30% or 40%. According to this assay, a functional homologous variant (ii) or functional fragment (iii) of natural exon sequence (i) is one which allows at least 50% of the expression improvement provided by the natural sequence.

Additionally or alternatively, exon sequence may be tested in the Comparative Immunogenicity Assay below. Test exon sequence is swapped into the base vector in place of the exon sequence already present. Functional exon sequence provides antibody titres that are at least as high as or higher than those achieved by the base vector with at least one, preferably two antigens. Preferably antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBsAg, HSV 2gD and flu-M2 antigens. According to this assay, a functional homologous variant (ii) or functional fragment (iii) of natural exon sequence (i) is one which allows the highest antibody titres achieved by the natural sequence.

The chimeric promoter construct comprises heterologous intron (c) in place of the native intron A region of the hCMV major immediate early gene. An intron is a non-coding sequence which is spliced from the hnRNA transcribed from a gene. A heterologous intron is one which is not present in the coding sequence in nature.

The heterologous intron (c) replaces wholly or partially, native intron A of the hCMV major immediate early gene. Typically the native intron A is absent.

In general the heterologous intron (c) is 3' of exon sequence (b).

Typically the heterologous intron (c) comprises:

- (i) a natural intron;
- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

Heterologous intron (c) is in general a viral or eukaryotic intron. Preferably the intron is a mammalian intron, in particular a non-human intron, for example a rat or chicken intron. Preferably the intron is an intron A, for example, rat insulin intron A, chicken keratin intron A or chicken cardiac action intron A.

Typically intron (c) has a length of from about 50 nucleotides to about 1000 nucleotides, for instance from about 100 to about 500 nucleotides. The intron (c) may for example, comprise 50 to 500 nucleotides, such as up to 100, 200, 300 or 400 nucleotides. Preferably the intron comprises sequence found at about nucleotides 50 to 133 of native rat insulin intron A, or a homologue of this sequence.

Preferably heterologous intron (c) is capable of being spliced from an RNA transcript in a eukaryotic host cell. In general the intron comprises one or more of a donor sequence (such as GT), an acceptor sequence (such as AG), a 3' pyrimidine rich region and a branch point sequence. The pyrimidine rich region, if present, may include, for example at least 5, 8, 10 or more pyrimidines. Preferably the intron comprises at least a donor sequence, acceptor sequence and a branch point sequence. Typically in the chimeric construct, intron (c) comprises non-intron flanking sequences which are derived from exon sequences found on the intron/exon boundaries of the natural intron (i). The flanking exon sequence may be native exon sequence or a homologue of this sequences which retains substantially the same activity as the native

sequence, for example retains splicing function. Typically from 5 to 10, preferably from 7 to 10 bases of exon sequence are included at each end of the intron.

Intron (c) may be an artificial intron, provided that the intron is functional. For example, a recombinant or chimeric intron may be used. Such an intron may comprise sequence from more than one natural intron.

Typically intron (c) comprises sequences present in hCMV intron A which bind transcription factors or regulatory proteins or instead of any of these sequences, homologues of these sequences able to bind the same factors or proteins. Typically such sequences or their homologues are present in the intron (c) in the same order and/or substantially the same relative spacing as in hCMV intron A.

Intron (c) may comprise an homologous variant (ii) in which the sequence of the natural intron (i) has been modified to remove an internal restriction site. For example, an homologous variant of rat insulin intron A may be used in which an internal *NheI* site has been destroyed.

Preferably, intron (c) comprises:

- (i) SEQ ID No. 3;
- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

Intron sequence (c) may be obtained using standard cloning techniques. For example, rat insulin intron A sequence is available at GenBank J00748, chicken keratin intron A sequence at GenBank J00847 and chicken cardiac intron A sequence at GenBank X02212. Intron

sequence can be isolated from natural sources using primers based on known sequence. Sequence may be prepared synthetically. Variant sequencer may be obtained by mutagenesis.

Typically a functional intron sequence, for example a functional variant (ii) or a functional fragment (iii) is one which has substantially the same activity as, and/or complements the activity of a natural intron (i). In one embodiment the activity is splicing activity.

Intron (c) sequences may be tested for splicing activity using a routine splicing assay. In general a functional homologue (ii) or functional fragment (iii) will show at least 50%, for example 60%, 70%, 80%, 90% and up to 100% or more of the splicing efficiency of the natural intron (i) in the assay.

In general the heterologous intron sequence will, when present in the construct of the invention, enhance expression. Typically, a variant (ii) or fragment (iii) intron will cause comparable enhancement to a natural intron (i).

Functionality of potential intron sequence (c) can be tested using the Comparative Expression Assay below. The heterologous intron is swapped into the base vector. Generally heterologous intron sequence is functional if the addition of the sequence increases expression in at least one, but preferably two reference cell types by 25% or more, compared to the base vector. Typically the reference cells are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. The increase in expression may be at least 35%, 45%, 55% or more. According to this assay, a functional variant (i) or functional fragment (ii) of a natural intron sequence (i) is one which allows greater than 50% of the expression improvement achieved by the natural sequence.

A heterologous intron (c) sequence may, additionally or alternatively be tested for functionality using the Comparative Immunogenicity Assay below. Intron (c) sequence is added to the base vector. A functional intron (c) sequence provides antibody titres that are higher than those achieved by the base vector with at least one, preferably two antigens. Preferably, the antibody titres are at least 5 or 10%, for instance 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBsAg, HSV2gD and Flu-M2 antigens. According to this assay, a functional variant (ii) or functional fragment (iii) of a natural intron sequence (i) is one which allows the highest antibody titres achieved by the natural sequence.

Suitable heterologous intron sequence can be obtained using standard cloning techniques. For example, rat insulin intron A sequence is available at GenBank J00748, chicken keratin intron A at GenBank J00847, and chicken cardiac actin intron A at X02212. Intron sequence can be isolated from native sources using primers based on the known sequence data. Suitable sequence may also be prepared synthetically.

The component sequences (a), (b) and (c) may be provided suitably linked together to form a chimeric promoter using standard cloning or molecular biology techniques. Preferably intron sequence (c) is provided 3' of exon sequence (b). The chimeric promoter construct is linked to a cloning site, in such a way that the promoter will effect the expression of a coding sequence inserted in the site, when the proper enzymes are present. Suitable cloning sites, including multi-cloning sites are known in the art, e.g the pUC19, pBC SK, pBluescript II KS, cDNA3.1, pSP72, pGEM 7Z multicloning site.

Typically, a nucleic acid for insertion (or inserted) in the cloning site encodes a therapeutically relevant polypeptide. It is preferred that the coding sequence is suitable for use in nucleic acid immunisation or gene therapy. The nucleic acid insert may thus comprise a sequence capable of providing immunity, for example an immunogenic sequence that elicits a humoral and/or cellular immune response when delivered to a subject. Alternatively, the nucleic acid may comprise one or more genes encoding a therapeutic polypeptide e.g a protein defective or missing from a target cell genome or a non-native protein having a desired biological or therapeutic effect (e.g., an antiviral function). For the treatment of genetic disorders, functional genes corresponding to genes known to be deficient in the particular disorder can be administered to a subject. Preferably the nucleic acid is DNA.

Suitable nucleic acids for insertion include those used for the treatment of inflammatory diseases, autoimmune, chronic and infectious diseases, including such disorders as AIDS, cancer, neurological diseases, cardiovascular disease, hypercholesterolemia; various blood disorders including various anemias, thalassemia and hemophilia; genetic defects such as cystic fibrosis, Gaucher's Disease, adenosine deaminase (ADA) deficiency, emphysema, etc.

For example, in methods for the treatment of solid tumors, genes encoding toxic peptides (i.e., chemotherapeutic agents such as ricin, diphtheria toxin and cobra venom factor), tumor suppressor genes such as p53, genes coding for mRNA sequences which are antisense to transforming oncogenes, antineoplastic peptides such as tumor necrosis factor (TNF) and other cytokines, or transdominant negative mutants of transforming oncogenes, can be inserted for expression at or near the tumor site.

Similarly, nucleic acids coding for polypeptides known to display antiviral and/or antibacterial activity, or stimulate the host's immune system, can also be included. The nucleic acid may encode one of the various cytokines (or functional fragments thereof), such as the interleukins, interferons and colony stimulating factors. The nucleic acid may encode an antigen for the treatment or prevention of a number of conditions including but not limited to cancer, allergies, toxicity and infection by a pathogen such as, but not limited to, fungus, viruses including Human Papilloma Viruses (HPV), HIV, HSV2/HSV1, influenza virus (types A, B and C), Polio virus, RSV virus, Rhinoviruses, Rotaviruses, Hepatitis A virus, Norwalk Virus Group, Enteroviruses, Astroviruses, Measles virus, Par Influenza virus, Mumps virus, Varicella-Zoster virus, Cytomegalovirus, Epstein-Barr virus, Adenoviruses, Rubella virus, Human T-cell Lymphoma type I virus (HTLV-I), Hepatitis B virus (HBV), Hepatitis C virus (HCV), Hepatitis D virus, Pox virus, Marburg and Ebola; bacteria including *M.tuberculosis*, *Chlamydia*, *N.gonorrhoeae*, *Shigella*, *Salmonella*, *Vibrio Cholera*, *Treponema pallidua*, *Pseudomonas*, *Bordetella pertussis*, *Brucella*, *Franciscella tulorensis*, *Helicobacter pylori*, *Leptospria interrogans*, *Legionella pneumophila*, *Yersinia pestis*, *Streptococcus (types A and B)*, *Pneumococcus*, *Meningococcus*, *Hemophilus influenza (type b)*, *Toxoplasma gondii*, *Compylobacteriosis*, *Moraxella catarrhalis*, *Donovanosis*, and *Actinomyces*; fungal pathogens including Candidiasis and Aspergillosis; parasitic pathogens including Taenia, Flukes, Roundworms, Amebiasis, Giardiasis, Cryptosporidium, Schistosoma, Pneumocystis carinii, Trichomoniasis and Trichinosis. The nucleic acid may also be used to provide a suitable immune response against numerous veterinary diseases, such as Foot and Mouth diseases, Coronavirus, *Pasteurella multocida*, *Helicobacter*, *Strongylus vulgaris*, *Actinobacillus pleuropneumonia*, Bovine viral diarrhea virus (BVDV),

Klebsiella pneumoniae, *E. Coli*, *Bordetella pertussis*, *Bordetella parapertussis* and *Bordetella brochiseptica*. Thus in one aspect, the nucleic acid constructs of the present invention may find use in a vaccine.

The nucleic acid for insertion into the cloning site may comprise a polyadenylation (polyA) sequence. Such a polyA sequence is generally native to the coding sequence. Alternatively, a heterologous polyA sequence may be provided in the nucleic acid construct of the invention. Typically the polyA sequence will be provided downstream of the cloning site, such that it is operatively linked to a coding sequence inserted in the cloning site. Any suitable polyA sequence may be included in the construct using standard cloning techniques. Such polyA sequences are known in the art.

The poly A sequence may be:

- (i) a natural poly A sequence;
- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

The natural poly A sequence (i) may be, for example a rabbit β globin gene poly A, Human Papilloma Virus (HPV) early or late gene poly A, HSV-2gB gene poly A, a simian CMV immediate early gene poly A or HSV gD late gene poly A.

Preferably the natural poly A sequence (i) is selected from the group consisting of SEQ ID No. 10 (GenBank K03256), SEQ ID No. 11 (GenBank M16019), SEQ ID No. 12 (GenBank Z80699) and SEQ ID No. 13 (GenBank K02718).

In general, a functional polyA sequence is one which retains polyadenylation activity.

A poly A sequence may be tested for the ability to bring about polyadenylation of an RNA transcript using a routine expression assay. A functional homologous variant (ii) or functional fragment (iii) typically shows at least 50%, for example 60%, 70% 80% or more of the poly A activity of the natural poly A sequence in the assay.

Generally the poly A sequence will, when present in the construct of the invention, enhance expression, typically causing comparable enhancement to the natural poly A sequence (i) mentioned above.

A poly A sequence may also be assayed for functionality using the Comparative Expression Assay below. A test poly A region is swapped into the base vector in place of the RBGpA. A test poly A is considered functional if the poly A does not abrogate expression but preferably increases expression in at least one but preferably two reference cell types, compared to the base vector. Preferably there is an increase in expression of at least 5%, 10%, 20%, 30%, 40% or 50% or more. Preferred cell types are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. According to the assay, an homologous variant (ii) or fragment (iii) is functional if it allows greater than 50% of the expression improvement achieved by the natural poly A sequence (i).

Alternatively, or additionally, poly A sequences may be tested for activity in the Comparative Immunogenicity Assay below. Poly A sequence is swapped into the base vector in place of RBGpA. A functional poly A sequence provides antibody titres that are at least as high as or higher than those achieved by the base vector with at least one, preferably two antigens. Preferably the antibody titres are at least

5%, 10%, for instance 20%, 30% or 40% higher than those achieved with the base vector. Preferred antigens are HBsAg, HSV2gD and Flu-M2 antigens. An homologous variant (ii) or fragment (iii) is functional if it allows the highest antibody titres achieved by the natural poly A sequence (i).

The nucleic acid construct may comprise additional control sequences which influence expression of a coding sequence inserted in the cloning site. The construct may include a non-translated leader sequence. The sequence is provided in the construct in operable linkage with the chimeric promoter, and therefore also with a coding sequence inserted at the cloning site. The leader provides a translational start site for expression of an inserted coding sequence and typically includes a Kozak sequence.

Typically the untranslated leader sequence comprises:

- (i) a natural untranslated leader sequence;
- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

In general the natural sequence (i) is a eukaryotic sequence or a viral sequence, in particular, of a virus which infects a eukaryote. Preferably the natural sequence (i) is HBV or HSV sequence, for example HBV preS2 antigen sequence, HBV e-antigen sequence, or HSV type 2gD antigen sequence. Particularly preferably, (i) is selected from the group consisting of SEQ ID No. 5, SEQ ID No. 6 and SEQ ID No. 7.

Typically the leader sequence comprises sequences present in (i) which bind transcription components or regulatory proteins, or

homologues of these sequences which are able to bind the same components or proteins. Typically such sequences or their homologues are present in the leader sequence in the same order and/or substantially the same relative spacing as in (i). In general the leader sequence comprises a translational start site for expression of an inserted coding sequence. Typically the leader sequence includes a Kozak sequence.

In general the untranslated leader sequence has a length of from about 10 to about 200 nucleotides, for example from about 15 to 150 nucleotides, preferably 15 to about 130 nucleotides. Leader sequences comprising, for example, 15, 50, 75 or 100 nucleotides may be used.

Generally a functional untranslated leader sequence is one which is able to provide a translational start site for expression of a coding sequence in operable linkage with the leader sequence. Typically a functional variant (ii) or fragment (iii) has substantially the same activity as and/or complements the activity of the natural sequence (i), usually in facilitating or enhancing expression of a coding sequence in operable linkage with the sequence.

A variant (ii) or fragment (iii) may be tested for activity as an untranslated leader sequence relative to natural leader sequence using standard protocols. For example, expression vectors may be prepared comprising a natural leader sequence (i) operably linked to its native coding sequence, and expression monitored in suitable host cells e.g. mammalian HEK 293T, CHO, HeLa, BHK, 3T3 or COS cells. Test constructs may be prepared in which the natural leader sequence is replaced by an homologous variant or fragment and expression is monitored again in the same host cells. In general, a variant (ii) or fragment (iii) provides at least 50%, such as 60%, 70%, 80%, 90% or 100% or more of the expression provided by the natural sequence.

A potential leader sequence can also be tested for utility in the Comparative Expression Assay below. A test leader sequence is swapped into the base vector in place of the HBV preS2 5'UTR. A functional leader sequence does not abrogate expression but preferably increases expression in at least one but preferably two reference cell types, compared to the base vector. In general expression is increased by at least 5%, 10%, 20%, 30%, 40% or 50%. Preferred cell types are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. According to the assay, an homologous variant (ii) or fragment (iii) is functional if it allows greater than 50% of the expression improvement achieved by the natural leader sequence.

Alternatively, or additionally, a leader sequence may be tested for activity in the Comparative Immunogenicity Assay below. A leader sequence is swapped into the base vector in place of HBV preS2 5'UTR. A functional leader sequence provides antibody titres that are at least as high as or higher than those achieved by the base vector, with at least one preferably two antigens. Preferably the antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBsAg, HSV 2gD and Flu-M2 antigens. An homologous variant (ii) or fragment (iii) is functional if it allows the highest antibody titres allowed by the natural leader sequence (i).

Suitable leader sequence can be obtained using standard protocols. For example, HBV preS2 antigen sequence, HBV e-antigen sequence and HSV type 2gD antigen sequence is available at GenBank M54923, M54923 and Z86099 respectively. Primers can be designed based on this known sequence and used to isolate homologous sequences. Leader sequences may be synthesised based on known sequences.

The nucleic acid construct may comprise an enhancer sequence. An enhancer sequence is typically provided 3' of the cloning site, in operable linkage with both the chimeric promoter and an inserted coding sequence, and acts to increase transcription of the inserted sequence.

In general the enhancer comprises:

- (i) a natural enhancer;
- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

The enhancer sequence generally comprises from about 50 to about 850 nucleotides, for example from about 75 to about 500 nucleotides. Enhancers of about 100, 200, 300 or 400 nucleotides may be used.

Typically (i) is a eukaryotic or viral enhancer, in particular, of a virus which infects eukaryotes. Usually such enhancers occur in the 3' untranslated region (3'UTR) of a gene. Preferably (i) is an HBV or a CMV enhancer, for example an HBs Ag 3' UTR or a simian CMV immediate early gene 3' UTR. Preferably (i) comprises SEQ ID No. 8 or SEQ ID No. 9.

In general, the enhancer in the construct comprises sequences found in (i) which bind transcription components or regulatory proteins, for example transcription factors, or homologues of these sequences which bind the same components or proteins. Preferably these sequences are present in the enhancer in the same order and/or substantially the same relative spacing as in (i).

Generally a functional enhancer is one which enhances or increases expression of a polynucleotide, for example, a coding sequence, which is operably linked to the enhancer sequence. Typically a functional homologous variant (ii) or fragment (iii) has substantially the same activity (for example, enhancement of expression) as and/or complements the activity of the natural enhancer (i).

Enhancer activity may be assayed using an enhancer trap assay. Protocols are known in the art. A functional homologous variant (ii) or fragment (iii) preferably provides at least 50% of the enhancer activity shown by the natural enhancer in such as assay. Typically the activity is at least 60%, 70%, 80%, 90, 100% or more of the activity of the natural enhancer. In general, a functional variant (ii) or fragment (iii) is able to complement the activity of the natural enhancer (i) in the assay.

Enhancer utility may also be tested using the Comparative Expression Assay set out below. A test 3'UTR sequence is swapped into the base vector. A 3'UTR has utility if it does not abrogate expression but preferably increases expression in at least one but preferably two reference cell types compared to the base vectors in the assay. Preferably expression is increased by at least 5%, 10%, 20%, 30%, 40% or 50%. Preferred cell types are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. According to this assay, an homologous variant (ii) or fragment (iii) is functional if it allows greater than 50% of the expression improvement achieved by the natural enhancer sequence (i).

Additionally or alternatively, enhancer sequences may be tested for activity in the Comparative Immunogenicity Assay below. A 3'UTR is swapped into the base vector. A functional enhancer sequence

provides antibody titres that are at least as high as or higher than those achieved by the base vector with at least one, preferably two antigens. Preferably the antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBs Ag, HSV2gD and Flu-M2 antigens. An homologous variant (ii) or fragment (iii) is functional if it allows the highest antibody titre allowed by the natural enhancer sequence (i).

Suitable enhancer sequence can be obtained using standard cloning methods. For example, HBsAg 3'UTR sequence, or simian CMV immediate early gene 3'UTR sequence can be accessed at GenBank AF143308 and M16019. Primers can be designed based on this known sequence and used to isolate homologous sequences.

In a preferred embodiment, the nucleic acid construct comprises a heterologous polyA sequence, a heterologous leader sequence and a heterologous enhancer, all in operable linkage with the chimeric promoter, for efficient expression of an inserted coding sequence.

In a further aspect, the present invention also provides a nucleic acid construct comprising, or sometimes consisting essentially of:

- (i) a promoter sequence
- (ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen sequence; and
- (iii) a coding sequence operably linked to (i) and (ii)

wherein the coding sequence is heterologous to the non-translated leader sequence.

Typically the promoter sequence (i) is derived from a viral or eukaryotic promoter sequence. The promoter sequence may be a natural promoter sequence, a functional homologue of the natural sequence or a functional fragment of either. Suitable natural promoters include, for example, the hCMV immediate early promoter, Pseudorabies virus (PRV) promoter or Rous sarcoma virus (RSV) promoter. Preferably the natural promoter comprises SEQ ID NO: 52 or SEQ ID NO: 53.

An artificial promoter construct, such as the chimeric promoter described above, may be used, provided that the promoter is functional.

A functional promoter sequence is generally one which is able to cause (including initiate and regulate) transcription of an operably linked coding sequence in a suitable host cell.

A promoter sequence may be tested for promoter activity using a routine expression assay. A functional homologue or fragment of a natural promoter sequence typically provides at least 50%, for example, or least 60, 70, 80 or 90% of the expression provided by the natural sequence in such an assay.

The non-translated leader sequence (ii) is as described above. Suitable coding sequences (iii) include those already described in relation to the chimeric promoter construct. However, in the present aspect of the invention, the coding sequence is heterologous to the non-translated leader sequence. The present construct typically includes a poly A sequence, which as already described, may be native to the coding sequence, or provided as a heterologous poly A sequence in the construct. Suitable poly A sequences have already been described. The construct may additionally include an enhancer sequence 3' of the

coding sequence. Suitable enhancer sequences are described above in relation to the chimeric promoter construct.

In another aspect, the invention provides a nucleic acid construct comprising, or in some embodiments consisting essentially of:

- (i) a promoter sequence;
- (ii) a coding sequence operably linked to the promoter sequence (i) and;
- (iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence, and the coding sequence (ii) is heterologous to the enhancer sequence.

The construct may include a non-translated leader sequence such as the ones already described in relation to the chimeric promoter construct.

Typically the promoter sequence (i) is derived from a viral or eukaryotic promoter sequence. The promoter sequence may be a natural promoter sequence, a functional homologue of the natural sequence or a functional fragment of either. Suitable natural promoters include, for example, the hCMV immediate early promoter, Pseudorabies virus (PRV) promoter or Rous sarcoma virus (RSV) promoter. Preferably the natural promoter comprises SEQ ID NO: 52 or SEQ ID NO: 53.

An artificial promoter construct, such as the chimeric promoter described above, may be used, provided that the promoter is functional.

A functional promoter sequence is generally one which is able to cause (including initiate and regulate) transcription of an operably linked coding sequence in a suitable host cell.

A promoter sequence may be tested for promoter activity using a routine expression assay. Functional homologues or fragments of a natural promoter sequence typically provide at least 50%, for example, or least 60, 70, 80 or 90% or the expression provided by the natural sequence in such an assay.

Suitable coding sequences (ii) include those already mentioned in relation to the chimeric promoter construct. However, in the present aspect, the coding sequence is heterologous to the 3' enhancer sequence. The enhancer sequence (iii) of the construct is described above. The present construct also typically includes a poly A sequence. As in the case of the chimeric promoter construct, this poly A region may be native to the coding sequence (ii) or may be provided as a heterologous poly A component in the construct.

A construct according to any aspect of the present invention may comprise a signal peptide sequence. The signal peptide sequence is inserted in operable linkage with the promoter such that the signal peptide is expressed and facilitates secretion of a polypeptide encoded by coding sequence also in operable linkage with the promoter.

Typically a signal peptide sequence encodes a peptide of 10 to 30 amino acids for example 15 to 20 amino acids. Often the amino acids are predominantly hydrophobic. In a typical situation, a signal peptide targets a growing polypeptide chain bearing the signal peptide to the endoplasmic reticulum of the expressing cell. The signal peptide is

cleaved off in the endoplasmic reticulum, allow for secretion of the polypeptide via the Golgi apparatus.

A signal peptide for use in the invention may comprise:

- (i) a natural signal peptide sequence;
- (ii) a homologous variant of (i) which retains signal peptide activity; or
- (iii) a fragment of (i) or (ii) which retains signal peptide activity.

Sequence (i) may be for example human tissue plasminogen activator signal peptide (hTPAsp) (GenBank L00141), the aprotinin signal peptide (GenBank AAD13685), tobacco extensin signal peptide (GenBank JU0465), or chicken lysosyme signal peptide (GenBank AF410481).

A signal peptide, suitable for use in the present invention, is one which will enable the secretion of heterologous proteins. A functional signal peptide can be identified in an assay which compares the effect of a test signal peptide with the effect of a known signal peptide - e.g. human tissue plasminogen activator signal peptide (hTPAsP) - and with the effect of having no signal peptide. The Comparative Expression Assay set out below may be used but with the following modification. Secretion expression vectors are constructed containing the base vector with either the test signal peptide, hTPAsp or no signal peptide. Coding sequences for polypeptides devoid of their naturally occurring signal peptides are inserted into the vectors and the vectors transformed into reference host cells. Preferably cells are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. The cell media is analysed for polypeptide expression levels. A functional signal peptide enables polypeptide

secretion at a higher level than a vector lacking a signal peptide with at least one, preferably two polypeptides. Typically, secretion is 5% higher, or more preferably 10% higher or more, for example 20 or 50% higher or more. Typically, secretion levels are comparable to those obtained using hTPAsp.

Allowing secretion of encoded protein outside of an expressing cell may have a number of advantages, in particular where the protein is an antigen. For example, increased antigen secretion could allow greater antigen uptake and response by immune cells (macrophages, Langerhans cells, B-cells, T-cells etc), enable the ability of antigen to reach the bloodstream and signal cells (cytokines), enable an antigen to find cellular ligands and effect a function (antibodies, toxins such as cholera toxin, *E.coli* LT) and participate in normal cellular biochemical processes (cellular receptors).

A nucleic acid construct of the invention may be in the form of a plasmid expression vector. The vector may then include additional elements, such as an origin of replication, or selector genes. Such elements are known in the art and can be included using standard techniques. In one embodiment, the plasmid vector has the sequence in SEQ ID NO:14. Alternatively, the construct may be included in a viral vector construct.

A polynucleotide construct of the invention may be substantially free of or associated with cells or with cellular material. It may be in substantially isolated form, or it may be in substantially purified form, in which case it will generally comprise at least 90% e.g. at least 95%, 98% or 99% of the polynucleotide or dry mass in the preparation.

The present nucleic acid molecules may be delivered to suitable host cells, for expression of a polynucleotide in operable linkage with the promoter. Preferably the host cells are mammalian cells, in particular human cells. Suitable methods for delivery of nucleic acids to such cells are known in the art and include, for example, dextran mediated transfection, calcium phosphate precipitation, electroporation and direct microinjection into nuclei.

As described above, a nucleic acid coding sequence in a construct may encode a therapeutically relevant polypeptide. The present constructs may therefore be used for nucleic acid immunisation or gene therapy using standard gene delivery protocols. Suitable methods for gene delivery are known in the art, as discussed below. The nucleic acid molecules can be delivered either directly to a subject, or alternatively, delivered *ex vivo* to cells derived from the subject whereafter the cells are reimplanted in the subject.

For use in nucleic acid immunisation or gene therapy, the nucleic acid constructs may be formulated as conventional pharmaceutical preparations. This can be done using standard pharmaceutical formulation chemistries and methodologies, which are available to those skilled in the art. For example, compositions containing one or more nucleic acid sequences (e.g., present in a suitable vector form such as a DNA plasmid) can be combined with one or more pharmaceutically acceptable excipients or vehicles to provide a liquid preparation.

Auxiliary substances, such as wetting or emulsifying agents, pH buffering substances and the like, may be present in the excipient or vehicle. These excipients, vehicles and auxiliary substances are generally pharmaceutical agents which may be administered without undue toxicity and which, in the case of vaccine compositions will not induce

an immune response in the individual receiving the composition. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, polyethyleneglycol, hyaluronic acid, glycerol and ethanol. Pharmaceutically acceptable salts can also be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. It is also preferred, although not required, that the preparation will contain a pharmaceutically acceptable excipient that serves as a stabilizer, particularly for peptide, protein or other like molecules if they are to be included in the composition. Examples of suitable carriers that also act as stabilizers for peptides include, without limitation, pharmaceutical grades of dextrose, sucrose, lactose, trehalose, mannitol, sorbitol, inositol, dextran, and the like. Other suitable carriers include, again without limitation, starch, cellulose, sodium or calcium phosphates, citric acid, tartaric acid, glycine, high molecular weight polyethylene glycols (PEGs), and combination thereof. A thorough discussion of pharmaceutically acceptable excipients, vehicles and auxiliary substances is available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991), incorporated herein by reference.

Certain facilitators of nucleic acid uptake and/or expression ("transfection facilitating agents") can also be included in the compositions, for example, facilitators such as bupivacaine, cardiotoxin and sucrose, and transfection facilitating vehicles such as liposomal or lipid preparations that are routinely used to deliver nucleic acid molecules. Anionic and neutral liposomes are widely available and well known for delivering nucleic acid molecules (see, e.g., *Liposomes: A Practical Approach*, (1990) RPC New Ed., IRL Press). Cationic lipid preparations are also well known vehicles for use in delivery of nucleic

acid molecules. Suitable lipid preparations include DOTMA (N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride), available under the tradename Lipofectin™, and DOTAP (1,2-bis(oleyl)-3-(trimethylammonio)propane), see, e.g., Felgner *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:7413-7416; Malone *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:6077-6081; US Patent Nos 5,283,185 and 5,527,928, and International Publication Nos WO 90/11092, WO 91/15501 and WO 95/26356. These cationic lipids may preferably be used in association with a neutral lipid, for example DOPE (dioleoyl phosphatidylethanolamine). Still further transfection-facilitating compositions that can be added to the above lipid or liposome preparations include spermine derivatives (see, e.g., International Publication No. WO 93/18759) and membrane-permeabilizing compounds such as GALA, Gramicidine S and cationic bile salts (see, e.g., International Publication No. WO 93/19768).

Alternatively, the nucleic acid molecules of the present invention may be encapsulated, adsorbed to, or associated with, particulate carriers. Suitable particulate carriers include those derived from polymethyl methacrylate polymers, as well as PLG microparticles derived from poly(lactides) and poly(lactide-co-glycolides). See, e.g., Jeffery *et al.* (1993) *Pharm. Res.* 10:362-368. Other particulate systems and polymers can also be used, for example, polymers such as polylysine, polyarginine, polyornithine, spermine, spermidine, as well as conjugates of these molecules.

Once formulated the compositions can be delivered to a subject *in vivo* using a variety of known routes and techniques. For example, the liquid preparations can be provided as an injectable solution, suspension or emulsion and administered via parenteral, subcutaneous, intradermal,

intramuscular, intravenous injection using a conventional needle and syringe, or using a liquid jet injection system. Liquid preparations can also be administered topically to skin or mucosal tissue, or provided as a finely divided spray suitable for respiratory or pulmonary administration. Other modes of administration include oral administration, suppositories, and active or passive transdermal delivery techniques.

Alternatively, the compositions can be administered *ex vivo*, for example delivery and reimplantation of transformed cells into a subject are known (e.g., dextran-mediated transfection, calcium phosphate precipitation, electroporation, and direct microinjection of into nuclei).

The compositions are administered to a subject in an amount that is compatible with the dosage formulation and that will be prophylactically and/or therapeutically effective. An appropriate effective amount will fall in a relatively broad range but can be readily determined by one of skill in the art by routine trials. The "Physicians Desk Reference" and "Goodman and Gilman's The Pharmacological Basis of Therapeutics" are useful for the purpose of determining the amount needed. For example, it is generally expected that an effective dose of the polynucleotide will fall within a range of about 0.001 to 1000 μ g, more preferably 0.01 to 10.0 μ g.

In a preferred embodiment, the nucleic acid constructs are delivered to target cells using a particle-mediated delivery technique. Particle mediated methods for delivering nucleic acid preparations are known in the art.

Particles for particle mediated delivery may be formed by coating the present nucleic acid molecules onto carrier particles (e.g., core carriers) using a variety of techniques known in the art. Carrier particles

are selected from materials which have a suitable density in the range of particle sizes typically used for intracellular delivery from a particle-mediated delivery device. Typically carrier particles have a diameter of from 0.1 to 5 μ m, for example 0.5 to 3 μ m, preferably 1 to 2 μ m. The optimum carrier particle size will, of course, depend on the diameter of the target cells. Alternatively, colloidal gold particles can be used wherein the coated colloidal gold is administered (e.g., injected) into tissue (e.g., skin or muscle) and subsequently taken-up by immune-competent cells.

Usually carrier particles are selected from inert metals. The metals are inert in that they are not physiologically active. For the purposes of the invention, tungsten, gold, platinum and iridium carrier particles can be used. Tungsten and gold particles are preferred. Tungsten particles are readily available in average sizes of 0.5 to 2.0 Φ m in diameter. Although such particles have optimal density for use in particle acceleration delivery methods, and allow highly efficient coating with DNA, tungsten may potentially be toxic to certain cell types. Gold particles or microcrystalline gold (e.g., gold powder A1570, available from Engelhard Corp., East Newark, NJ) will also find use with the present methods. Gold particles provide uniformity in size (available from Alpha Chemicals in particle sizes of 1-3 Φ m, or available from Degussa, South Plainfield, NJ in a range of particle sizes including 0.95 Φ m) and reduced toxicity. Microcrystalline gold provides a diverse particle size distribution, typically in the range of 0.1-5 Φ m. However, the irregular surface area of microcrystalline gold provides for highly efficient coating with nucleic acids.

A number of methods are known and have been described for coating or precipitating DNA or RNA onto gold or tungsten particles.

Most such methods generally combine a predetermined amount of gold or tungsten with plasmid DNA, CaCl_2 and spermidine. The resulting solution is vortexed continually during the coating procedure to ensure uniformity of the reaction mixture. After precipitation of the nucleic acid, the coated particles can be transferred to suitable membranes and allowed to dry prior to use, coated onto surfaces of a sample module or cassette, or loaded into a delivery cassette for use in particular particle-mediated delivery instruments.

As an alternative, the polynucleotides of the invention can be formulated as a particulate composition. Formulation can be carried out using the above-described standard pharmaceutical formulation chemistries. For example, the polynucleotides can be combined with one or more pharmaceutically acceptable excipients or vehicles to provide a suitable composition. The formulated compositions are then prepared as particles using standard techniques such as by simple evaporation (air drying), vacuum drying, spray drying, freeze drying (lyophilisation), spray-freeze drying, spray coating, precipitation, supercritical fluid particle formulation, and the like. If desired, the resultant particles can be densified using the techniques described in commonly owned International Publication No. WO 97/48485, incorporated herein by reference.

These methods can be used to obtain nucleic acid particles having a size ranging from about 0.01 to about 250 μm , preferably about 10 to about 150 μm , and most preferably about 20 to about 60 μm ; and a particle density ranging from about 0.1 to about 25 g/cm^3 , and a bulk density of about 0.5 to about 3.0 g/cm^3 , or greater.

Once formed, the particles comprising the nucleic acid molecules may be packaged in single unit dosages or multidose containers. Such

containers may comprise an hermetically sealed container enclosing a suitable amount of the particles. The particles can be packaged as a sterile formulation, and the hermetically sealed container can thus be designed to preserve the sterility of the formulation until use in delivery to a subject. The containers are preferably adapted for direct use in a particle mediated delivery device. Typically such containers take the form of capsules, foil pouches, sachets, cassettes and the like. The particle delivery devices can also be provided in a preloaded condition containing a suitable dosage of the particles. The preloaded device may then also be prepackaged in a hermetically sealed container.

The container in which the particles are packaged can further be labelled to identify the composition and provide relevant dosage information. In addition, the container can be labelled with a notice in the form prescribed by a governmental agency, for example, the Food and Drug Administration, wherein the notice indicates approval by the agency under Federal Law of the manufacture, use or sale of the nucleic acid preparation contained therein for human administration.

Particle acceleration devices, suitable for particle-mediated delivery are known in the art. For example, current gene gun devices employ an explosive, electric or gaseous discharge to propel coated carrier particles towards target cells. The coated carrier particles can be releasably attached to a movable carrier sheet, or removably attached to a surface along which a gas stream passes, lifting the particles from the surface and accelerating them toward the target. An example of a gaseous discharge device is described in U.S. Patent No. 5,204,253. An explosive-type device is described in U.S. Patent No. 4,945,050. One example of an electric discharge apparatus suitable for use herein is described in U.S. Patent No. 5,120,657. Another electric discharge

apparatus is described in US Patent No 5,149,655. The disclosure of all of these patents is incorporated herein by reference in their entireties.

Particles may also be administered using a needleless syringe device, such as those described in U.S. Patent No. 5,630,796 to Bellhouse *et al* ("the PowderJect® needleless syringe device") and in International Publication Nos. WO 94/24263, WO 96/04947, WO 96/12513 and WO 96/20022, all of which are incorporated herein by reference.

Devices such as the one described in US Patent No. 5,630,796 may be provided as a pen-shaped instrument containing, in linear order moving from top to bottom, a gas cylinder, a particle cassette or package, and a supersonic nozzle with an associated silencer medium. The particles are provided within a suitable container, e.g. a cassette formed by two rupturable polymer membranes that are heat-sealed to a washer-shaped spacer to form a self-contained sealed unit. Membrane materials can be selected to achieve a specific mode of opening and burst pressure that dictate the conditions at which the supersonic flow is initiated.

In operation, the device is actuated to release the compressed gas from the cylinder into an expansion chamber within the device. The released gas contacts the particle cassette and, when sufficient pressure is built up, suddenly breaches the cassette membranes sweeping the particles into the supersonic nozzle for subsequent delivery. The nozzle is designed to achieve a specific gas velocity and flow pattern to deliver a quantity of particles to a target surface of predefined area. The silencer is used to attenuate the noise produced by the supersonic gas flow.

The delivery system described in International Publication No. WO 96/20022 also uses the energy of a compressed gas source to accelerate and deliver powdered compositions. However, it is distinguished from the system of US Patent No. 5,630,796 in its use of a shock wave instead of gas flow to accelerate the particles. More particularly, an instantaneous pressure rise provided by a shock wave generated behind a flexible dome strikes the back of the dome, causing a sudden eversion of the flexible dome in the direction of a target surface. This sudden eversion catapults a powdered composition (which is located on the outside of the dome) at a sufficient velocity, thus momentum, to penetrate target tissue, e.g., oral mucosal tissue. The powdered composition is released at the point of full dome eversion. The dome also serves to completely contain the high-pressure gas flow which therefore does not come into contact with the tissue. Because the gas is not released during this delivery operation, the system is inherently quiet. This design can be used in other enclosed or other wise sensitive applications for example, to deliver particles to minimally invasive surgical sites.

Particles may be delivered *in vivo* directly to a subject, or *ex vivo* to cells taken from a subject, the transformed cells then being reimplanted in the subject. For *in vivo* delivery, particle injection is typically subcutaneously, epidermally, intradermally, intramucosally (e.g. nasally, rectally and/or vaginally), intraperitoneally, intravenously, orally or intramuscularly. Preferably, delivery is to terminally differentiated cells; however, the particles can also be delivered to non-differentiated, or partially differentiated cells such as stem cells of blood and skin fibroblasts. Most preferably, delivery is to skin epidermal cells.

The particles are administered to a subject in a manner compatible with the dosage formulation and in an amount that will be prophylactically and/or therapeutically effective. A "therapeutically effective amount" of the present particulate compositions will be sufficient to bring about treatment or prevention of disease or condition symptoms, and will fall in a relatively broad range that can be determined by routine trials. Generally the particles are delivered in an amount of from 0.001 to 1000 μ g, more preferably 0.01 to 10.0 μ g of nucleic acid per dose. However, the exact amount necessary will vary depending on the age and general condition of the individual being treated and the particular nucleotide sequence selected, as well as other factors. An appropriate effective amount can be readily determined through clinical testing. The "Physicians Desk Reference" and "Goodman and Gilman's The Pharmacological Basis of Therapeutics" are useful for the purpose of determining the amount needed.

Assays

Comparative Expression Assay

A suitable test for element utility determines the effect the element has on expression of a polypeptide. The basis of comparison for testing utility of the elements is a 'base vector', generally (unless otherwise noted) a plasmid with a hCMV promoter, hCMV exon 1, 9 bases of hCMV exon 2, the 5'UTR from HBV preS2 and the rabbit- β globin polyadenylation region, positioned to drive expression of a coding sequence. Typically, the base vector is pJV7384, pJV 7401, pJV 7450 or pJV 7533. Heterologous introns and 3'UTRs are added, or promoter sequences, exons, 5'UTRs and polyA sites are swapped into the base vectors to create test expression vectors. Thus functional variants or fragments can be tested.

The base vectors and test vectors are transformed into suitable host cells and the cells analysed for polypeptide expression levels. Preferably mammalian host cells are used. Suitable cells include mammalian HEK 293T, CHO, HeLa, BHK, 3T3 or COS cells.

Typically, a functional element causes expression which is comparable to the base vector, for example at least the same as or greater. Preferably expression is tested in more than one cell type and with more than one coding sequence.

Suitable experimental protocols are provided, for example, in Examples 1 to 13 below.

Comparative Immunogenicity Assay

Where the polypeptide to be expressed is an antigen, a further test may be carried out to identify functional or particularly preferred construct elements. In the assay, the effect of an element on immune response is determined after delivery of an expression vector to a test organism. Antibody levels against the antigen are the easiest way to judge immune response. Groups of mice are vaccinated with the base vectors or test vectors constructed as above. Sera is collected after an appropriate amount of time and analyzed for antibody levels.

This experiment is performed twice, and the antibody levels from all the groups in both experiments are plotted. Functional elements will give rise to at least as high as or higher antibody titres in both experiments for a particular antigen than the base vector. Preferably, the result will be seen with more than one antigen to demonstrate the breadth of utility of the element(s) in that expression panel.

Suitable experimental protocols are provided for example, in Example 14 below.

C. Experimental

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (*e.g.*, amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

Methods

Standard PCR Conditions

The standard PCR conditions used for the construction of vectors were as follows: 1x PCR core buffer w/ 1.5mM MgCl₂ (Promega Corporation, Madison, WI), 0.400μM each of each primer, 200μM of each dNTP (USB, Inc, Cleveland, OH) , 2.5μ Taq polymerase (Promega Corporation, Madison, WI), 1.0 ng template DNA, water to 100μl, and a mineral oil (Aldrich Chemical, Inc, Milwaukee, WI) overlay. The PTC-200 thermocycler (MJ Research, Inc, Waltham, MA) was programmed to run the following routine: 4'@95EC, 30 cycles of (1'@95EC/ 1'15"@55EC/ 1'@72EC), 10'@72EC, 4EC hold). The amplification products were removed from the PCR reaction by using the QIAquickâPCR Purification Kit (Qiagen Inc, Valencia, CA) prior to cutting with restriction enzymes (New England Biolabs, Beverly, MA).

All PCR products were sequenced after cloning to ensure fidelity of the amplification.

Example 1. Construction of hepatitis B virus surface antigen (HBsAg) vector panels

A number of plasmid expression vectors were constructed for expression of HBsAg.

Starting materials

(i) pWRG7128 (Roy, M, *et.al. Vaccine* (2001) 19: 764-778), which contains the hCMV immediate early promoter sequence, the first exon, first intron, and a partial second exon of the hCMV major

immediate early gene, the HBsAg coding sequence with flanking regions (HBV preS2 5'UTR derived sequence and 3' posttranscriptional response element) and the bovine growth hormone polyadenylation region (BGHpA)

(ii) pJV7284, a derivative of pWRG7128 that exchanges the rabbit β globin polyadenylation region (RBGpA) for BGHpA.

(a) pJV7384 (CMV(no intron), HBV preS2 5'UTR and RBGpA)

pWRG7128 was PCR'd with JF93 (SEQ ID NO:15) and F110 (SEQ ID NO:16) using standard conditions and cut with *Sa*I and *Bam*H1 to isolate an insert fragment containing the CMV promoter, exon 1 and part of the exon 2 sequence. pAM6 (ATCC, Mannassas, VA) was cut with *Bam*H1 and *Bst*X1 to isolate an insert fragment that contained the 5'-UTR of HBsAg, and roughly 70% of the HBsAg coding region. pJV7284 was cut with *Sa*I and *Bst*X1 to generate a vector fragment into which the two insert fragments were ligated, resulting in pJV7293.

pWRG7128 was PCR'd with primers GW1 (SEQ ID NO:17) and JF254 (SEQ ID NO:18) and cut with *Bst*X1 and *Bgl*II to isolate an insert fragment that contained the 3'-end of the HBsAg coding region.

pJV7293 was cut with *Bst*X1 and *Bgl*II to generate a vector fragment into which the insert fragment was ligated, resulting in vector pJV7384.

(b) pJV7382 (CMV(no intron), HBsAg 3'UTR, HBV preS2 5'UTR and RBGpA)

pJV7293 was cut with *Xho*I and *Xba*I to generate an insert fragment containing the CMV promoter/exons and the 5'-UTR with 5'-end of the HBsAg coding sequence. pWRG7128 was cut with *Xba*I and *Bcl*I to generate an insert fragment containing the majority of the HBsAg

coding sequence and the 3'-UTR. pJV7284 was cut with *Xho*1 and *Bgl*2 to generate a vector fragment into which the two insert fragments were ligated, resulting in pJV7382.

(c) pJV7389 (CMV (RIA), HBsAg 3'UTR, HBV preS2 5'UTR and RBGpA)

The rat insulin intron A (RIA) was PCR'd out of plasmid p5'rlns (unknown origin) with primers GW150 (SEQ ID NO:19) and JF255 (SEQ ID NO:20). The PCR product was cut with *Bam*H1 and inserted into *Bam*H1 linearized pJV7382, resulting in pJV7389.

(d) pJV7387 (CMV(RIA), HBV preS2 5'UTR and RBGpA)

pJV7384 was cut with *Bst*X1 and *Eco*R1 to generate an insert fragment containing the 3' end of the HBsAg coding region and RBGpA. pJV7389 was cut with *Bst*X1 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7387.

Example 2. Construction of herpes simplex virus glycoprotein D antigen (HSVgD) vector panels

A number of plasmid expression vectors were constructed for expression of HSVgD.

Starting materials

(a) pJV7334, a derivative of pWRG7284 (pJV 7284) that replaces the HBsAg coding sequence with an in-frame *Nhe*1 directly downstream of the ATG start codon, followed by a stuffer fragment with a *Bam*H1 immediately 5' of the HBV Enh

(b) pWRG7202 , a derivative of pGem3Z (Promega) with a stuffer fragment that allows the fusion of a coding sequence to the human tissue plasminogen activator (TPA) signal peptide downstream of a *Nhe*1 site.

(a) pJV7392(CMV(native intron), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

The coding region for HSV2 gD was PCR'd out of a viral DNA stock (Advanced Biotech, Inc, Columbia, MD) using primers DS1 (SEQ ID NO:21) and DA1 (SEQ ID NO:22) and was cut with *Nhe*1 and *Eco*R1 to generate an insert fragment. pWRG7202 was cut with *Nhe*1 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7391.

pJV7391 was cut with *Nhe*1 and *Bgl*/2 to generate an insert fragment containing the HSV2 gD coding sequence. pJV7334 was cut with *Nhe*1 and *Bam*H1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7392. This vector consists of the following expression elements: the hCMV immediate early promoter sequence, the first exon, first intron, and a partial second exon of the hCMV major immediate early gene, the 5'-UTR from HBsAg, the coding sequence for HSV2 gD gene, the 3'-UTR from HBsAg, and RBGpA.

(b) pJV7399 (CMV(no intron), HBsAg 3'UTR HBsAg 5'UTR and RBGpA)

An intronless version of pJV7392 was constructed as follows. pJV7384 was cut with *Hind*3 and *Nde*1 to isolate an insert fragment containing the 5' ends of the kanamycin resistance gene and the CMV

promoter. pJV7384 was cut with *Nde*1 and *Ssp*1 to isolate an insert fragment containing the 3' end of the CMV promoter, the CMV exon1/2 and 5' end of the 5'-UTR from HBsAg. These insert fragments were inserted into pJV7392 from which the *Hind*3-*Ssp*1 fragment was removed, resulting in pJV7399.

(c) pJV7400 (CMV(RIA), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

A RIA version of pJV7392 was constructed as follows. pJV7384 was cut with *Hind*3 and *Nde*1 to isolate an insert fragment containing the 5' ends of the kanamycin resistance gene and CMV promoter. pJV7387 was cut with *Nde*1 and *Ssp*1 to isolate an insert fragment containing the 3' end of the CMV promoter, the CMV exon1/2(partial), RIA, and 5' end of the 5'-UTR from HBsAg. These insert fragments were inserted into pJV7392 from which the *Hind*3-*Ssp*1 fragment was removed, resulting in pJV7400.

(d) pJV7401 (CMV(no intron), HBsAg 5'UTR and RBGpA)

A 3'-UTR-less version of pJV7399 was constructed as follows. pJV7391 was cut with *Bsp*120I and *Bgl*2 to isolate an insert fragment containing the 3' end of the HSV2 gD gene. pJV7284 was cut with *Bgl*2 and *Eco*R1 to isolate the RBGpA signal. These insert fragments were inserted into pJV7399 from which the *Bsp*120I-*Eco*R1 fragment was removed, resulting in pJV7401.

(e) pJV7402 (CMV(RIA), HBsAg 5'UTR and RBGpA)

A 3'-UTR-less version of pJV7400 was constructed as follows. pJV7391 was cut with *Bsp*120I and *Bgl*2 to isolate an insert fragment containing the 3' end of the HSV2 gD gene. pJV7284 was cut with *Bgl*2

and *EcoR1* to isolate the RBGpA signal. These insert fragments were ligated into pJV7400 from which the *Bsp120I-EcoR1* fragment was removed, resulting in pJV7402.

Example 3. Construction of Flu M2 antigen vector panels

(a) pJV7450 (CMV(no intron), HBsAg 5'UTR and RBGpA)

A coding region for Flu M2 was PCR'd out of plasmid pFL-M2 (Joel Haynes, PJV) using primers JF301 (SEQ ID NO:23) and JF302 (SEQ ID NO:24) and was cut with *Nhe1* and *Bgl2* to generate an insert fragment. pJV7401 was cut with *Nhe1* and *Bgl2* to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7450.

(b) pJV7452 (CMV(no intron), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

A 3'UTR fragment was PCR'd out of pJV7389 with primers JF84 (SEQ ID NO:25) and JF225 (SEQ ID NO:26) was cut with *Bsp120I*, filled with T4 DNA polymerase, and linkerized with *Bgl2* linkers (cat# 1036, New England Biolabs). The fragment was then cut with *Bgl2* and *EcoR1* to isolate an insert fragment containing the 3'UTR of HBsAg and the RBGpA region. pJV7450 was cut with *Bgl2* and *EcoR1* to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7452

(c) pJV7458 (CMV(RIA) HBsAg 5' UTR and RBGpA)

A version of pJV7450 containing the RIA was constructed as follows: pJV7389 was cut with *BamH1* to isolate a RIA containing insert

fragment. pJV7450 was cut with *Bam*H1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7458.

(d) pJV7468 (CMV(RIA), HBsAg 3'UTR ,HBsAg 5'UTR and RBGpA)

A version of pJV7458 containing the 3'UTR of HBsAg was constructed as follows: pJV7452 was cut with *Bgl*/2 and *Eco*R1 to produce an insert fragment containing the HBsAg 3'UTR and RBGpA. pJV7458 was cut with *Bgl*/2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7468.

Example 4. Construction of Beta-gal vector panels

(a) pJV7488 (CMV(no intron), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

CMV-beta (Clontech) was PCRed with primers JF335 (SEQ ID NO:27) and JF336 (SEQ ID NO:28) and cut with *Nhe*1 and *Bgl*/2 to isolate an insert fragment coding for beta-galactosidase. pJV7452 was cut with *Nhe*1 and *Bgl*/2 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7488.

(b) pJV7533 (CMV(no intron), HBsAg 5'UTR and RBGpA)

pJV7450 was cut with *Bgl*/2 and *Eco*R1 to isolate an insert fragment containing the RBGpA. pJV7488 was cut with *Bgl*/2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7533.

(c) pJV7551(CMV(RIA/NheI), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

pJV7530 (see Example 5) was cut with *Xho*I and *Bam*HI to isolate an insert fragment containing the CMV promoter through RIA. pJV7488 was cut with *Xho*I and *Bam*HI to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7551.

(d) pJV7552(CMV(RIA/NheI), HBsAg 5'UTR and RBGpA)

pJV7530 was cut with *Xho*I and *Bam*HI to isolate an insert fragment containing the CMV promoter through RIA. pJV7533 was cut with *Xho*I and *Bam*HI to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7552.

Example 5. Construction of pJV Expression (pJV7563)

(a) pJV7496

pJV7389 was PCR'd with primers JF357 (SEQ ID NO:29) and JF365 (SEQ ID NO: 30), treated with T4 DNA polymerase to blunt the ends, and cut with *Sa*I to isolate an insert fragment coding for kanamycin resistance. pJV7389 was cut with *Ava*I, treated with T4 DNA polymerase to blunt the ends, and cut with *Sa*I to isolate a vector fragment into which the insert fragment was ligated, resulting in pJV7496.

(b) pJV7530

pJV7389 was PCR'd with primers JF393 (SEQ ID NO:31) and JF406 (SEQ ID NO:32) and cut with *Bgl*II and *Bam*HI to isolate an insert fragment containing the RIA devoid of an internal *Nhe*I site. pJV7496

was cut with *Bam*H1 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7530.

(c) pJV7549

pJV7468 was cut with *Bam*H1 and *Eco*R5 to isolate an insert fragment containing M2 and part of the HBV 3'ENH. pJV7530 was cut with *Bam*H1 and *Eco*R5 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7549.

(d) pJV7563

Primers JF256 (SEQ ID NO:33) and JF257 (SEQ ID NO:34) were annealed to prepare an insert fragment consisting of a multiple cloning site. pJV7549 was cut Nhe1 and Bgl2 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7563. A pJV7563 plasmid map is provided in Figure 12. The base composition for the pJV7563 plasmid is provided in Figure 13. The components and their position in the plasmid pJV7563 are as follows:

1-44 Transposon 903 sequences

45-860 Kanamycin resistance coding sequence from
Transposon 903

861-896 Transposon 903 sequences

897-902 Sal1 site

903-1587 CMV promoter

1588-1718 untranslated leader sequence from the immediate-early
gene of CMV

1719-1724 Fusion of BamH1 and BglII restriction enzymes

1725-1857 Rat insulin intron A

1858-1863 BamH1 site

1864-1984 HBV surface antigen 5'- untranslated leader

1985-1993 Synthetic start codon/ Nhe1 cloning site

1994-2011 Synthetic cloning sites

2012-2544 HBV enhancer

2545-2555 Old vector sequence. No hits against NCBI databases

2556-2686 Rabbit beta-globin polyadenylation region

2687-3759 pUC19 vector sequence

**Example 6. Construction of Signal Peptide Expression Panels using
Human Secreted Alkaline Phosphatase (SEAP) and Human IgG Fc
Fragment (hFc) as Model Antigens**

(i) pJV7507 (hTPAsp and SEAP)

pSEAP-Basic (Clontech) was PCR'd with primers JF320 (SEQ ID NO:35) and JF321 (SEQ ID NO:36) then cut with *Nhe1* and *Bgl2* to isolate an insert fragment consisting of the human SEAP fragment. pJV7079 (Macklin, et.al.) was cut with *Nhe1* and *Bgl2* to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7507.

(ii) pJV7508 (hTPAsp and hFc)

Human DNA was PCR'd with primers JF386 (SEQ ID NO:37) and FcAS (SEQ ID NO:38) then cut with *Nhe*1 and *Bgl*2 to isolate an insert fragment consisting of the human IgG Fc fragment. pJV7079 was cut with *Nhe*1 and *Bgl*2 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7508.

(iii) Preparation of Aprotinin Signal Peptide Coding Sequence

Synthetic oligo JF354 (SEQ ID NO:39) was PCR'd with primers JF355 (SEQ ID NO:40) and JF356 (SEQ ID NO:41) to generate the coding sequence for the aprotinin signal peptide.

(iv) Preparation of Tobacco Extensin Signal Peptide Coding Sequence

Synthetic oligo JF348 (SEQ ID NO:42) was PCR'd with primers JF349 (SEQ ID NO:43) and JF350 (SEQ ID NO:44) to generate the coding sequence for the tobacco extensin signal peptide.

(v) Preparation of Chicken Lysozyme Signal Peptide Coding Sequence

Synthetic oligo JF351 (SEQ ID NO:45) was PCR'd with primers JF352 (SEQ ID NO:46) and JF353 (SEQ ID NO:47) to generate the coding sequence for the chicken lysozyme signal peptide.

(a) Flu M2 antigen signal peptide panels

pJV7499 (CMV(no intron), HBsAg5'UTR, RBGpA, aprotinin s.p.)

pJV7497 (CMV(no intron), HBsAg5'UTR, RBGpA, tobacco extensin s.p.)

PJV7500 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken lysosyme s.p.).

Coding sequences for the signal peptides were cut with *Spe1* and *Nhe1* to isolate insert fragments. pJV7450 was cut with *Nhe1* to prepare a vector fragment into which the insert fragments were ligated, resulting in pJV7499 (aprotinin), pJV7497 (tobacco extensin), and pJV7500 (chicken lysozyme).

(b) SEAP signal peptide panels

pJV7513 (CMV(no intron), HBsAg5'UTR, RBGpA, aprotinin s.p.)

pJV7512 (CMV(no intron), HBsAg5'UTR, RBGpA, tobacco extensin sp.)

pJV7510 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken lysosyme s.p.)

pJV7499, 7497, and 7500 were cut with *Xho1* and *Nhe1* to isolate an insert fragment consisting of the CMV promoter through the signal peptide coding sequence of the plasmids. PJV7507 was cut with *Xho1* and *Nhe1* to prepare a vector fragment into which the insert fragments were ligated, resulting in pJV7513 (aprotinin), pJV7512 (tobacco extensin), and pJV7510 (chicken lysozyme).

(c) hFc signal peptide panels

pJV7524 (CMV(no intron), HBsAg5'UTR, RBGpA, aprotinin s.p.)

pJV7525 (CMV(no intron), HBsAg5'UTR, RBGpA, tobacco extensin s.p.)

pJV7526 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken lysosyme signal peptide)

pJV7499, 7497, and 7500 were cut with *Xho*1 and *Nhe*1 to isolate an insert fragment consisting of the CMV promoter through the signal peptide coding sequence of the plasmids. pJV7508 was cut with *Xho*1 and *Nhe*1 to prepare a vector fragment into which the insert fragments were ligated, resulting in pJV7524 (aprotinin), pJV7525 (tobacco extensin), and pJV7526 (chicken lysozyme).

Example 7. Construction of Human Secreted Alkaline Phosphatase (SEAP) panels

(a) pJV7531 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken lysosyme s.p.)

pJV7510 was cut with *Sa*/1 and *Bg*/2 to isolate an insert fragment containing the CMV promoter through lysozyme signal peptide. pJV7450 was cut with *Sa*/1 and *Bg*/2 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7531.

(b) pJV7554 (CMV(RIA/*Nhe*l), HBsAg5'UTR, RBGpA, chicken lysosyme s.p.)

pJV7530 was cut with *Xho*1 and *Bam*H1 to isolate an insert fragment containing the CMV promoter through RIA. pJV7531 was cut with *Xho*1 and *Bam*H1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7554.

(c) pJV7568 (CMV(no intron), HBsAg 3'UTR, HBsAg5'UTR, RBGpA, chicken lysosyme s.p.)

pJV7563 was cut with *Bgl*2 and *Eco*R1 to isolate an insert fragment containing the HBV 3'-UTR and RBGpA. pJV7531 was cut with *Bgl*2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7568.

(d) pJV7572 (CMV(RIA/*Nhe*l), HBsAg 3'UTR, HBsAg5'UTR, RBGpA, chicken lysosyme s.p.)

pJV7563 was cut with *Bgl*2 and *Eco*R1 to isolate an insert fragment containing the HBV 3'UTR and RBGpA. pJV7554 was cut with *Bgl*2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7572.

Example 8. Construction of Beta-gal and HBsAg Vectors Using the Chicken Keratin and Chicken Cardiac Actin Introns

(a) pJV7557 (Beta-gal, CMV(cA intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

Chicken DNA was PCR'd with primers JF430 (SEQ ID NO:48) and JF442 (SEQ ID NO:49) and cut with *Bgl*2 and *Bam*H1 to isolate an insert fragment consisting of the intron and flanking exon sequences from chicken cardiac actin. pJV7488 was cut with *Bam*H1 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7557.

(b) pJV7558 (Beta-gal, CMV(cK intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

Chicken DNA was PCR'd with primers JF421 (SEQ ID NO:50) and JF444 (SEQ ID NO:51) and cut with *Bgl*II and *Bam*HI to isolate an insert fragment consisting of the intron and flanking exon sequences from the chicken keratin gene. pJV7488 was cut with *Bam*HI to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7558.

(c) pJV7578 (HBsAg, CMV(cA intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

pJV7557 was cut with *Sal*I and *Bam*HI to isolate an insert fragment consisting of the CMV promoter through intron regions. pJV7496 was cut with *Sal*I and *Bam*HI to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7558.

(d) pJV7579 (HBsAg, CMV(cK intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

pJV7558 was cut with *Sal*I and *Bam*HI to isolate an insert fragment consisting of the CMV promoter through intron regions. pJV7496 was cut with *Sal*I and *Bam*HI to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7579.

Example 9. *In-Vitro* Analysis of Antigen Expression by HBsAg

Vector Panels

On day one, SSC15 (ATCC) or B16 (origin unknown, versions available at ATCC) cells were plated on 6 well tissue culture plates at 20-40% confluency, and allowed to grow overnight in an incubator. The host cells were propagated in media recommended by ATCC.

On day two, the transfection reaction was performed. For each vector to be tested, 20 μ l of Lipofectin[®] reagent (Life Technologies Inc, Grand Island, NY) was added to 180 μ l of Optimem[®] media (Life Technologies, Grand Island, NY), and allowed to incubate at room temperature for 45 minutes. For each vector to be tested, 2 μ g of vector was mixed into 200 μ l of Optimem[®] at 40 minutes. At 45 minutes, the vector and Lipofectin[®] solutions were mixed together and allowed to sit at room temperature for an additional 10 minutes. During this final incubation, the plated host cells were removed from the incubator and washed twice with Optimem[®] media. At 10 minutes, 1.6ml of Optimem[®] was added to the Lipofectin[®] / vector mix, and 1ml of the resultant mix was added to each of two cell wells. The host cells were returned to the incubator and allowed to sit undisturbed for 5 hours, at which point the Lipofectin[®] / vector mix was removed and replaced by standard cell maintenance media.

At 18 to 24 hours after the media change, from 50 to 100 μ l of cell maintenance media was removed from the tissue culture plates and analyzed for antigen expression by placing the samples into reaction vessels provided in the AUSZYME[®] Monoclonal Diagnostic Kit (Abbott Laboratories, Abbott Park, IL). The volume of the test samples was brought to a volume of 200 μ l with PBS, then 50 μ l of conjugate and a reaction bead were added to each sample. The vessel is incubated for 80

minutes at 40EC, after which the wells were washed clean of all liquid reaction components. The beads were transferred to new tubes after which 300 μ l of color development buffer was added. At 30 minutes, the color development reaction was stopped by the addition of 1M sulfuric acid, and the absorbance of the reaction was measured at 490nm. The data shown in Figure 1 is the average absorbance readings of the duplicate wells from two experiments.

As shown in Figure 1, the addition of RIA, the HBV 3'UTR or both elements to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of HBsAg in SSC15 cells. As shown in Figure 2, the addition of either the chicken keratin or the chicken cardiac actin intron to a base vector (CMV promoter, exon, HBV 3'UTR and polyadenylation region) increased expression of HBsAg in SSC15 cells.

Example 10. *In-Vitro* Analysis of Antigen Expression by Beta-gal Vector Panels

SSC-15 or B16 host cells were transfected as described in Example 9.

At eighteen to forty hours after the media change, the media supernatants were removed and the cells were washed with PBS. After removal of the wash, the cells were lysed by incubating the cells in 500 μ l lysis buffer (50mM NaPO₄, 0.1% Triton X-100, pH 7) for 5 minutes, followed by physically scraping the cells off the plastic dish. The lysates were microfuged for two minutes to remove cell debris, and 10 to 25 μ l of the cleared lysate were added to 500 μ l of reaction buffer (80ug/ml o-nitrophenyl galactopyranoside, 50mM NaPO₄, pH 7) and incubated at 37EC for 10 to 20 minutes. The reaction was stopped by the addition of 500 μ l of 1M Na₂CO₃ and read at 405 nm. Data is

presented as the ratio of the expression of enhanced (containing an intron, HBVenh, or both) vector to a base vector.

The addition of RIA, the HBV 3'UTR or both elements to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of beta-gal in both cell lines. The results for SSC15 cells are shown in Figure 3. Addition of either the chicken keratin or the chicken cardiac actin intron to a base vector (CMV promoter, exon, the HBV 3'UTR and polyadenylation region) increased the expression of beta-gal in both cell lines. Results for B16 cells are shown in Figure 2.

Example 11. In-Vitro Analysis of Antigen Expression by HSV gD Vector Panel

SSC15 or B16 host cells were transfected as described in Example 9. Eighteen hours post transfection, plates were placed on ice for 15 minutes. Each well was then washed with 2 ml of PBS (Biowhittaker, Walkerville, MD). Cells were fixed with 0.05% gluteraldehyde (Polysciences Inc, Warrington, PA) diluted in PBS and incubated for 30 minutes at room temperature. All subsequent incubations lasted 1 hour at room temperature and washes between each incubation were as stated above. The plates were blocked with 2 ml of 5% dry milk (Bio Rad Laboratories, Melville, NY) in PBS. Incubations with 1 ml of a 1:1000 dilution of anti-gD monoclonal (ABI, Columbia, MD) in 2% dry milk / PBS / 0.05% Tween-20 (Sigma, St. Louis, MO) and 1 ml of a 1:2500 dilution of goat anti-mouse HRP (KPL, Gaithersburg, MD) in PBS / 0.1% Tween-20 followed. Color was developed using 1 ml of TMB microwell substrate (BioFX, Owings Mills, MD). The reactions were stopped with 1M H₂SO₄, the liquid was transferred to plastic cuvettes and the optical density read at 450nm. Data is presented as the ratio of the expression

of enhanced (containing an intron, HBVenh, or both) vector to a base vector.

Addition of RIA with or without the HBV 3'UTR to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of HSV gD in both cell lines. Results for SC15 cells are shown in Figure 4.

Example 12. *In-Vitro* Analysis of Antigen Expression by SEAP Vector Panels

SSC15 or B16 host cells were transfected as described in Example 9. At eighteen to forty hours after the media change, the media supernatants were removed and heated at 70EC for 30 minutes. 10 to 25 μ l of the heat-inactivated supernatants were incubated for 5 minutes with 1/10th volume of 100mM L-homoarginine. 500 μ l of alkaline phosphatase reaction buffer (cat # 172-1063, Bio-Rad, prepared according to instructions) were added to the lysates and incubated at 37EC for 10 to 20 minutes. The reaction was stopped by the addition of 500 μ l of 1M NaOH and read at 405 nm. Data is presented as the ratio of the expression of enhanced (containing an intron, HBVenh, or both) vector to a base vector, or the ratio of the expression of the experimental signal peptides to the human TPA signal peptide vector.

As shown in Figure 5, the addition of RIA, the HBV 3'UTR or both elements to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of SEAP in B16 cells. Unexpectedly, only the addition of the HBV 3'UTR to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of SEAP in SSC15 cells.

Addition of signal peptides from either bovine aprotinin, chicken lysozyme, or tobacco extensin to the N-terminus of mature SEAP allowed for efficient secretion of SEAP into cell media supernatants of both cell lines. Results for B16 cells are shown in Figure 6.

Example 13. In-Vitro Analysis of Antigen Expression by Human IgG Fc Fragment Signal Peptide Panel

SSC15 or B16 host cells were transfected as described in Example 9. The media supernatants were removed from eighteen to forty hours after the media change.

ELISA plates (Costar) were incubated overnight at 4°C with 100 µl of goat anti-human IgG (Sigma #13382, 1/1000 dilution in carbonate coating buffer) per well. All subsequent incubations lasted 1 hour at room temperature with washes (10mM Tris, 150mM NaCl, 0.1% Brij-35, pH8.0) between each incubation. The wells were then blocked with 100µl of 5% dry in PBS, followed by incubation with serially diluted media supernatants in dilution buffer (2% dry milk, PBS, 0.05% Tween-20). This was followed by incubation with 100 µl of goat anti-human IgG/HRP (Sigma #A6029, 1/5000 dilution in dilution buffer) per well, followed by color development using 100µl of TMB microwell substrate. The reactions were stopped with 100µl of 1M H₂SO₄, and read at 450 nm. Data is presented as the ratio of the expression of the experimental signal peptides to the human TPA signal peptide vector.

Addition of signal peptides from either bovine aprotinin, chicken lysozyme, or tobacco extensin to the N-terminus of the human Fc fragment allowed for efficient secretion of hFc into cell media supernatants of both cell lines. Results for B16 cells are shown in Figure 6.

Example 14. Use of the HBsAg, HSVgD and Flu-M2 plasmid expression vectors for immunisation of mice

a. Preparation of cartridges of immunisation

For each plasmid to be tested, 25mg of 2 micron gold powder was weighed into a microfuge tube. After the addition of a 250 μ l aliquot of 50 mM spermidine (Aldrich Chemical, Inc, Milwaukee, WI), the tube was vortexed and briefly sonicated. The gold was microfuged out, and the spermidine replaced by a fresh 100 μ l aliquot. The gold was resuspended by vortexing, after which 25 μ g of DNA was added to the tube and mixed. While the tube was lightly vortexed, 100 μ l of 10% CaCl (Fujisawa USA, Inc, Deerfield, IL) was added to precipitate the DNA onto the gold beads. The precipitation reaction was allowed to proceed for 10 minutes on the benchtop, after which the gold was collected by a brief microfuge spin and washed three times with absolute ethanol (Spectrum Quality Products, Inc, Gardena, CA) to remove excess precipitation reagents. The washed gold/DNA complex was then resuspended in 3.6 ml of 0.05 mg/ml polyvinylpyrrolidone (360KD, Spectrum Quality Products, Inc, Gardena, CA) in absolute ethanol. This slurry was then injected into a Tefzelâ tube (McMaster-Carr, Chicago, IL) located in a tube turner (PowderJect Vaccines) which coats the inside of the Tefzelâ tube with the gold/DNA complex. After the tube turning procedure was completed, the tube was cut into 0.5" "shots" of vaccine which were loaded into the XR1 device (PowderJect Vaccines) for delivery to the mice.

b. Vaccination Procedure

Four to six week old mice were anesthetized with a mixture of Ketasetâ (Fort Dodge) and Rompunâ (Bayer). The bellies were shaved with a pair of electric clippers to remove hair, and two non-overlapping "shots" of vaccine were delivered via the XR1 device (450psi) to the shaved area. The animals were returned to their cages and bled at six weeks post vaccination. Balb/c mice were used to evaluate the HBsAg expression vectors, and Swiss Webster mice were used to evaluate the HSV-gD and Flu M2 expression vectors.

Analysis of Sera for Anti-HBsAg Antibodies

At six weeks, blood samples were harvested from vaccinated animals. A volume of serum isolated from these samples was placed into wells of a reaction vessel supplied with the AUSAB® EIA Diagnostic Kit (Abbott Laboratories, Abbott Park, IL). The volume of sera added depended upon the antibody titer of the sample, and the sample was diluted with sample dilution buffer to fall within values obtainable with a quantification panel. 200µl from each vial of the AUSAB® Quantification Panel (Abbott Laboratories, Abbott Park, IL) was added to wells of the reaction vessel. To each well a bead was added, after which the vessel was sealed and incubated for two hours at 40EC. The wells were then washed of all liquid reaction components. To each washed well was added 200µl of conjugate mix, after which the vessel was sealed and incubated for two hours at 40EC. The wells were then washed of all liquid reaction components. The beads were transferred to new tubes after which 300µl of color development buffer was added. At 30 minutes, the color development reaction was stopped by the addition of 1M sulfuric acid, and the absorbance of the reactions was measured at 490nm in a Quantum II® spectrophotometer (Abbott Laboratories,

Abbott Park, IL) . This spectrophotometer calculates the antibody levels of a sample by comparing the absorbance of the sample with a standard curve generated with the quantification panel. These antibody levels were then corrected for dilution factors. The data shown in Figure 7 are the geometric mean titers of all animals vaccinated with a particular vector.

Analysis of Sera for Anti-Flu M2 Antigen Antibodies

96-well Costar medium-binding ELISA plates (Fisher Scientific, Pittsburgh, PA) were coated with a synthetic Flu M2 peptide (QCB / Biosource, Hopkinton, MA) at a concentration of 1ug/ml in PBS (Biowhittaker, Walkerville, MD) and incubated overnight at 4EC. The plates were washed three times with 10mM Tris (Sigma, St. Louis, MO) / 150mM NaCl (Fisher Scientific) / 0.1 % Brij-35 (Sigma), then blocked with 5% dry milk (Bio Rad Laboratories, Melville, NY) in PBS for 1 hour at room temperature. All subsequent incubations were at room temperature for one hour and washes between each incubation were as stated above. Sample mouse sera, a standard (high titer, anti-M2 mouse sera) and a negative control (anti-HBsAg mouse sera) were diluted in 2% dry milk / PBS / 0.05% Tween-20 (Sigma) and incubated in the ELISA plates. Goat anti-mouse IgG (H + L) biotin conjugated antibody (Southern Biotechnology Associate, Birmingham, AL) diluted 1:8000 in 2% dry milk / PBS / 0.05% Tween-20 and streptavidin-horseradish peroxidase conjugate (Southern Biotechnology) diluted 1:8000 in PBS / 0.1% Tween-20 followed. Color was developed using TMB substrate (BioFX, Owings Mills, MD). The reactions were stopped with 1M H₂SO₄ and the plates read at 450nm with an Emax precision microplate reader (Molecular Devices, Sunnyvale, CA). SoftMax Pro 4.1 software (Molecular Devices) was used to calculate endpoint titers using a four-

parameter analysis. Titers were normalized to the standard serum, which had a pre-determined titer, to minimize assay-to-assay and plate-to-plate variation. Results are shown in Figure 7.

Analysis of Sera for Anti-HSV gD Antigen Antibodies

96-well Costar medium-binding ELISA plates (Fisher Scientific, Pittsburgh, PA) were coated with HSV gD (Viral Therapeutics, Ithaca, NY) protein at a concentration of 1ug/ml in PBS (Biowhittaker, Walkerville, MD) and incubated overnight at 4EC. The plates were washed three times with 10mM Tris (Sigma, St. Louis, MO) / 150mM NaCl (Fisher Scientific) / 0.1% Brij-35 (Sigma), then blocked with 5% dry milk (Bio Rad Laboratories, Melville, NY) in PBS for 1 hour at room temperature. All subsequent incubations were at room temperature for one hour and washes between each incubation were as stated above. Sample mouse sera, a standard (high titer, anti-gD mouse sera) and a negative control (anti HBsAg mouse sera) were diluted in 2% dry milk / PBS / 0.05% Tween-20 (Sigma) and incubated in the ELISA plates. Goat anti-mouse IgG (H + L) biotin conjugated antibody (Southern Biotechnology Associate, Birmingham, AL) diluted 1:8000 in 2% dry milk / PBS / 0.05% Tween-20 and streptavidin-horseradish peroxidase conjugate (Southern Biotechnology) diluted 1:8000 in PBS / 0.1% Tween-20 followed. Color was developed using TMB substrate (BioFX, Owings Mills, MD). The reactions were stopped with 1M H₂SO₄ and the plates read at 450nm with an Emax precision microplate reader (Molecular Devices, Sunnyvale, CA). SoftMax Pro 4.1 software (Molecular Devices) was used to calculate endpoint titers using a four-parameter analysis. Titers were normalized to the standard serum, which had a pre-determined titer, to minimize assay-to-assay and plate-to-plate variation. Results are shown in Figure 7.

Accordingly, novel nucleic acid constructs, compositions comprising these constructs, and nucleic acid immunization techniques using these constructs have been described. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

Brief Description of the Sequences

SEQ ID NO:1 is hCMV immediate early promoter sequence
(GenBank #M60321, X17403)

SEQ ID NO:2 is sequence from exons 1 and 2 of the hCMV major
immediate early gene (GenBank #M60321, X17403)

SEQ ID NO:3 is rat insulin intron A sequence (GenBank #J00748)

SEQ ID NO:4 is the sequence of a chimeric promoter according to
the present invention

SEQ ID NO:5 is a leader sequence from HBV preS2 antigen 5' UTR
sequence (GenBank #M54923)

SEQ ID NO:6 is a leader sequence from HSV type 2 gD 5'UTR
sequence (GenBank #Z86099)

SEQ ID NO:7 is a leader sequence from HBV e antigen 5'UTR
sequence (GenBank #M54923)

SEQ ID NO:8 is HBVenh 3'UTR sequence (GenBank #AF143308)

SEQ ID NO:9 is simian immediate early gene 3'UTR sequence
(GenBank #M16019)

SEQ ID NO:10 is rabbit β globin poly A sequence (GenBank
#K03256)

SEQ ID NO:11 is simian sCMV immediate early gene poly A
sequence (GenBank #M16019)

SEQ ID NO:12 is HSV2 gB gene poly A sequence (GenBank
#Z86099)

SEQ ID NO:13 is HPV16 early gene poly A sequence (GenBank #K02718)

SEQ ID NO:14 is the sequence of the pJV expression vector

SEQ ID NO:15 is PCR primer JF93

SEQ ID NO:16 is PCR primer F110

SEQ ID NO:17 is PCR primer GW1

SEQ ID NO:18 is PCR primer JF254

SEQ ID NO:19 is PCR primer GW150

SEQ ID NO:20 is PCR primer JF255

SEQ ID NO:21 is PCR primer DS1

SEQ ID NO:22 is PCR primer DA1

SEQ ID NO:23 is PCR primer JF301

SEQ ID NO:24 is PCR primer JF302

SEQ ID NO:25 is PCR primer JF84

SEQ ID NO:26 is PCR primer JF225

SEQ ID NO:27 is PCR primer JF335

SEQ ID NO:28 is PCR primer JF336

SEQ ID NO:29 is PCR primer JF357

SEQ ID NO:30 is PCR primer JF365

SEQ ID NO:31 is PCR primer JF393

SEQ ID NO:32 is PCR primer JF406

SEQ ID NO:33 is PCR primer JF256

SEQ ID NO:34 is PCR primer JF257

SEQ ID NO:35 is PCR primer JF320

SEQ ID NO:36 is PCR primer JF321

SEQ ID NO:37 is PCR primer JF386

SEQ ID NO:38 is PCR primer FcAS

SEQ ID NO:39 is oligonucleotide JF354

SEQ ID NO:40 is PCR primer JF355

SEQ ID NO:41 is PCR primer JF356

SEQ ID NO:42 is oligonucleotide JF348

SEQ ID NO:43 is PCR primer JF349

SEQ ID NO:44 is PCR primer JF350

SEQ ID NO:45 is oligonucleotide JF351

SEQ ID NO:46 is PCR primer JF352

SEQ ID NO:47 is PCR primer JF353

SEQ ID NO:48 is PCR primer JF430

SEQ ID NO:49 is PCR primer JF442

Atty. Dkt. No.: 036481/0141

SEQ ID NO:50 is PCR primer JF421

SEQ ID NO:51 is PCR primer JF444

SEQ ID NO:52 is Pseudorabies virus (PRV) promoter sequence.

SEQ ID NO:53 is Rous sarcoma virus (RSV) promoter sequence.

SEQUENCE LISTING

<110> JAMES FULLER

<120> NUCLEIC ACID CONSTRUCTS

<130> APF 33 (X.13149 GCW)

<160> 53

<170> PatentIn version 3.1

<210> 1

<211> 685

<212> DNA

<213> Human cytomegalovirus

<400> 1

aatattgget attggccatt gcatacgttg tatctatata ataatatgta catttatatt	60
ggctcatgtc caatatgacc gccatgttga cattgattat tgactagtta ttaatagtaa	120
tcaattacgg ggtcattagt tcatagccca tatatggagt tccgcgttac ataacttacg	180
gtaaatggcc cgcctggetg accgcccac gaccccgcc cattgacgtc aataatgacg	240
tatgttccca tagtaacgcc aatagggact ttccattgac gtcaatgggt ggagtattta	300
cggtaaactg cccacttggc agtacatcaa gtgtatcata tgccaagtcc gccccctatt	360
gacgtcaatg acggtaaatg gccgcctgg cattatgcc agtacatgac cttacgggac	420
tttcotactt ggcagtacat ctacgtatta gtcacgcta ttaccatggt gatgcggttt	480
tggcagtaca ccaatgggcy tggatagcgg ttgactcac ggggatttcc aagtctccac	540
cccattgacg tcaatgggag ttgttttgg caccaaaac aacgggactt tccaaaatgt	600
cgtaataacc cgcgccgtt gacgcaaatg ggcggtaggc gtgtacggtg ggaggtatat	660
ataagcagag ctcgtttagt gaacc	685

<210> 2

<211> 131

<212> DNA

<213> Human cytomegalovirus

<400> 2

gtcagatcgc ctggagacgc catccacgct gttttgacct ccatagaaga caccgggacc	60
gatccagcct ccgcgccggg gaacggtgca ttggaacgcy gattccccgt gccaaagatg	120
actcaccgtc c	131

<210> 3
 <211> 135
 <212> DNA
 <213> Rattus rattus

<400> 3
 atcagcaagc aggtatgtac tctccagggt gggcctggct tccccagtca agactccagg 60
 gatttgaggg acgotgtggg ctcttctctt acatgtacct ttgctagcc tcaaccctga 120
 ctatcttcca ggtca 135

<210> 4
 <211> 955
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Chimeric promoter sequence

<400> 4
 aatattggct attggccatt gcatacgttg tatctatato ataatatgta catttatatt 60
 ggctcatgtc caatatgacc gccatgttga cattgattat tgactagtta ttaatagtaa 120
 tcaattaagg ggtcattagt tcatagccca tatatggagt tccgcgttac ataacttacg 180
 gtaaattggc cgcctggctg accgcccac gaccccgcc cattgacgtc aataatgacg 240
 tatgttccca tagtaacgcc aatagggact ttccattgac gtcaatgggt ggagtattta 300
 cggtaaactg cccacttggc agtacatcaa gtgtatcata tgccaagtcc gcccctatt 360
 gacgtcaatg acggtaaatg gcccgctgg cattatgcc agtacatgac cttacgggac 420
 tttcctactt ggcagtacat ctacgtatta gtcacgcta ttaccatggt gatgcggttt 480
 tggcagtaca ccaatggcg tggtatagcg tttgactcac ggggatttcc aagtctccac 540
 cccattgacg tcaatgggag tttgttttgg caccaaaac aacgggactt tccaaaatgt 600
 cgttaatacc ccgccccgtt gacgcaaatg ggcggtaggc gtgtacggtg ggaggtctat 660
 ataagcagag ctctgttagt gaaccgtcag atcgctgga gacgccatcc acgctgtttt 720
 gacctccata gaagacacgg ggaccgatcc agcctccgcg gccgggaacg gtgcattgga 780
 acgcggttc cccgtgccaa gactgactca cgtccggat ctcagcaagc aggtatgtac 840
 tctccagggt gggcctggct tccccagtca agactccagg gatttgaggg acgctgtggg 900
 ctcttctctt acatgtacct ttgcttgcc tcaaccctga ctatcttcca ggtca 955

<210> 5
 <211> 121
 <212> DNA

<213> Hepatitis B virus

<400> 5

cagagtcagg ggtotgtatt ttcctgctgg tggotccagt tcaggaacag taaacctgc	60
tccgaatatt gcctctcaca tctcgtcaat ctccgcgagg actggggacc ctgtgacgaa	120
c	121

<210> 6

<211> 57

<212> DNA

<213> Herpes simplex virus

<400> 6

ataagctgca ttgcgaacca ctagtcgcgcg ttttctgtgt gcacgcgta tcacggc	57
--	----

<210> 7

<211> 48

<212> DNA

<213> Hepatitis B virus

<400> 7

ctttgtacta ggaggctgta ggcataaatt ggtctgttca ccagcacc	48
--	----

<210> 8

<211> 533

<212> DNA

<213> Hepatitis B virus

<400> 8

taacaaaaca aaaagatggg gttattccct aaacttcatg ggttacgtaa ttggaagttg	60
ggggacattg ccacaagatc atattgtaca aaagatcaaa cactgtttta gaaaacttcc	120
tgtaaacagg cctattgatt ggaaagtatg tcaaaggatt gtgggtcttt tgggctttgc	180
tgctccattt acacaatgtg gatatcctgc cttaatgcct ttgtatgcat gtatacaage	240
taaacaggct ttcactttct cgccaactta caaggccttt ctaagtaaac agtacatgaa	300
cctttacccc gttgctcggc aacggcctgg tctgtgccaa gtgtttgctg acgcaacccc	360
cactggctgg ggottggcca taggccaatca ggcgatgcgt ggaacctttg tggctcctct	420
gcgatccat actgeggaaac tcctagccgc ttgttttgcct cgcagccggg ctggagcaaa	480
gctcatagga actgacaatt ctgtcgtcct ctccgggaaa tatacatcgt ttc	533

<210> 9
 <211> 158
 <212> DNA
 <213> Simian cytomegalovirus

<400> 9
 gtcagacaga cagacagtta tatgggctgg tccotataac totgccattg taaccccata 60
 tagccagaca gttagcattg catctattga tgatgtacta atgtattgta accccccta 120
 tgccattgtc taactgtact aatgtatgat attatacc 158

<210> 10
 <211> 131
 <212> DNA
 <213> Oryctolagus cuniculus

<400> 10
 gatctttttc cctctgcca aaattatggg gacatcatga agccccttga gcatctgact 60
 totggctaataa aaaggaaatt tattttcatt gcaatagtgt gttggaattt tttgtgtctc 120
 tcaactcggaa g 131

<210> 11
 <211> 204
 <212> DNA
 <213> Simian cytomegalovirus

<400> 11
 atatatactc tatgttatac tctatgatat acaatatata ctcatgaaca ctatgtactt 60
 ggtgtatgac tcattattgt ctgggaactg gttgggactt ggttggttgg gaagaatgtt 120
 gtgcctgtac ttgtgctgtg ctgtggatct caataaatgt gactatgttc aaaacactaa 180
 gtgccccgt gtcttcttta acta 204

<210> 12
 <211> 163
 <212> DNA
 <213> Herpes simplex virus 2

<400> 12
 gaagacgagc tctaaggag gggaggggag ctgggcttgt gtataaataa aaagacaccg 60
 atgttcaaaa atacacatga cttctggtat tggttttcct tggtttttat ttgggggggg 120

gggggcgtgt gactagaaaa acaaatgcag acatgtgcta acg

163

<210> 13

<211> 191

<212> DNA

<213> Human papillomavirus type 16

<400> 13

aattgttaca tataattggt gtataccata acttactatt tttcttttt tttttcata	60
tataattttt tttttgttt gtttgttgt ttttaataa actgttatta cttaacaatg	12
cgacacaaac gttctgcaaa acgcacaaaa cgtgcacgg ctacccaact ttataaaca	180
tgcaaacagg c	191

<210> 14

<211> 3759

<212> DNA

<213> Artificial sequence

<220>

<223> pJV expression vector

<220>

<221> Intron

<222> (1725)..(1857)

<223> Rat Ins IntA

<220>

<221> misc_feature

<222> (1)..(44)

<223> Tn903, pUC4K Remnants

<220>

<221> misc_feature

<222> (861)..(896)

<223> Tn903, pUC4K Remnants

<220>

<221> misc_feature

<222> (897)..(902)

<223> pUC19 MCS

<220>

<221> polyA_signal

<222> (2556)..(2686)

<223> rGLOB pA

<220>

<221> polyA_site

<222> (2647)..(2647)

<223> PolyA_Site_1

<220>

<221> promoter

<222> (903)..(1587)

<223> CMV Pro

<220>

<221> 3'UTR

<222> (2012)..(2544)

<223> HBVenh

<220>

<221> 5'UTR

<222> (1864)..(1984)

<223> 5'-UTR of HBV pre-S2

<220>

<221> misc_feature

<222> (1719)..(1724)

<223> Bam/Bgl fusion

<220>

<221> misc_feature

<222> (1985)..(1987)

<223> ATG-Nhe

<220>

<221> misc_feature

<222> (1988)..(2011)

<223> CDS insertion site

<220>

<221> misc_feature
<222> (2545)..(2555)
<223> unknown

<220>
<221> exon
<222> (1588)..(1718)
<223> CMV Exon 1/2

<220>
<221> misc_feature
<222> (2693)..(3759)
<223> pUC19

<220>
<221> misc_feature
<222> (45)..(860)
<223> KanR (Tn903) complement

<400> 14
ggcgtaatgc tctgccagtg ttacaaccaa ttaaccaatt ctgattagaa aaactcatcg 60
agcatcaaat gaaactgcaa tttattcata tcaggattat caataccata tttttgaaaa 120
agcogtttct gtaatgaagg agaaaactca ccgaggcagt tccataggat ggcaagatcc 180
tggatcggt ctgcgatcc gactcgtcca acatcaatac aacctattaa tttccctcg 240
tcaaaaataa ggttatcaag tgagaaatca ccatgagtga cgaactgaatc cggtgagaat 300
ggcaaaagct tatgcatttc tttccagact tgttcaacag gccagccatt acgctcgtca 360
tcaaaatcac tcgcatcaac caaacggtta ttcattcgtg attgogcctg agcgagacga 420
aatacgcgat cgctgttaaa aggacaatta caaacaggaa tcgaatgcaa cgggcgcagg 480
aacactgcca gcgcatcaac aatatattca cctgaatcag gatattcttc taatacctgg 540
aatgotgttt tcccggggat cgcagtggg agtaaccatg catcatcagg agtacggata 600
aatgcttga tggtcggaag aggcataaat tccgtcagcc agtttagtct gaccatctca 660
tctgtaacat cattggcaac gctaccttg ccatgtttca gaaacaactc tggcgcatcg 720
ggcttcccat acaatcgata gattgtcgca cctgattgcc cgacattatc gcgagcccat 780
ttatacccat ataaatcago atccatgttg gaatttaac gcggcctcga gcaagacgtt 840
tcccggtgaa tatggctcat aacacccctt gtattactgt ttatgtaagc agacaggtcg 900
acaatattgg ctattggcca ttgcatacgt tgtatctata tcataatatg tacatttata 960
ttggctcatg tccaatatga ccgccatgtt gacattgatt attgactagt tattaatagt 1020
aatcaattac ggggtcatta gttcatagcc catatatgga gttccgcgtt acataactta 1080
cggtaaatgg cccgcctggc tgacgcgcca acgacccccg cccattgacg tcaataatga 1140
cgtatgttcc catagtaacg ccaataggga ctttccattg acgtcaatgg gtggagtatt 1200
tacggtaaac tgccacttg gcagtacatc aagtgtatca tatgccaagt cggcccccta 1260

ttgacgtcaa tgacggtaaa tggcccgcct ggcattatgc ccagtacatg accttacggg 1320
 aactttcctac ttggcagtag atctacgtat tagtcatcgc tattaccatg gtgatgcggg 1380
 tttggcagta caccaatggg cgtggatagc ggtttgactc acggggattt ccaagtctcc 1440
 accccattga cgtcaatggg agtttgtttt ggcacaaaaa tcaacggggac tttccaaaat 1500
 gtcgtaataa ccccgccccc ttgacgcaaa tgggcggtag gcgtgtacgg tgggaggtct 1560
 atataagcag agctcgttta gtgaacc gtc aga tcg cct gga gac gcc atc cac 1614

Val Arg Ser Pro Gly Asp Ala Ile His

1

5

gct gtt ttg acc tcc ata gaa gac acc ggg acc gat cca gcc tcc gcg 1662
 Ala Val Leu Thr Ser Ile Glu Asp Thr Gly Thr Asp Pro Ala Ser Ala
 10 15 20 25

gcc ggg aac ggt gca ttg gaa cgc gga ttc ccc gtg cca aga gtg act 1710
 Ala Gly Asn Gly Ala Leu Glu Arg Gly Phe Pro Val Pro Arg Val Thr
 30 35 40

cac cgt cc ggatctcagc aagcaggtat gtactctcca ggggtggcct ggcttcccca 1768
 His Arg

gtcaagactc cagggatttg agggacgctg tgggtctctt tcttacatgt accttttgot 1828
 tgcctcaacc ctgactatct tccaggtcag gatcccagag tcaggggtct gtattttcct 1888
 gctggtggct ccagttcagg aacagtaaac cctgotccga atattgcctc tcacatctcg 1948
 tcaatctccg cagggactgg ggaccctgtg acgaacatgg ctacgaggcc cagatctggg 2008
 ccctaacaaa acaaaaagat ggggttatct cctaaacttc atgggttacg taattggaag 2068
 ttgggggaca ttgccacaag atcatattgt acaaaagatc aaacactgtt ttagaaaact 2128
 tctgttaaac aggcctattg attgaaaagt atgtcaaagg attgtgggtc ttttgggctt 2188
 tgetgtcca tttacacaat gtggatatcc tgccttaatg cctttgtatg catgtataca 2248
 agctaaacag gctttcaact totogccaac ttacaaggcc tttctaagta aacagtacat 2308
 gaacctttac cccgttgctc ggcaacggcc tggctctgtc caagtgttg ctgacgcaac 2368
 cccactggc tggggcttgg ccataggcca tcagcgcatg cgtggaacct ttgtggctcc 2428
 tetgcegatc catactgogg aactcctage cgcttgtttt gctcgcagcc ggtctggagc 2488
 aaagctcata ggaactgaca attctgtcgt cctctcggcg aaatatacat cgtttcgatc 2548
 tacgtatgat ctttttccct ctgccaaaaa ttatggggac atcatgaagc cccttgagca 2608
 tctgaactct ggctaataaa ggaatttat tttcattgca atagtgtgtt ggaatttttt 2668
 gtgtctctca ctcggaagga attctgcatt aatgaatcgg ccaacgcgcg gggagaggcg 2728
 gtttgcgat tgggcgctct tccgcttccct cgtcactga ctgctgcgc tcggctcgtc 2788
 ggctcggcg agcggatatc gctcactcaa aggcggtaat acggttatcc acagaatcag 2848
 gggataacgc aggaagaagc atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa 2908
 aggcgcgctt gctggcgctt ttccataggg tcgccccccc tgacgagcat cacaaaaatc 2968
 gacgctcaag tcagaggtgg cgaaaccgca caggactata aagataccag gcgtttcccc 3028

ctggaagctc cctcgtgcgc tctcctgttc cgaccctgcc gcttaccgga tacctgtccg	3088
cctttctccc ttcggaagc gtggcgcttt ctcatagctc acgctgtagg tatctcagtt	3148
cggtgtaggt cgttcgctcc aagctgggct gtgtgcacga acccccgtt cagcccgacc	3208
gctgcgcctt atccggtaac tatcgtcttg agtccaaccc ggtaagacac gacttatcgc	3268
cactggcagc agccactggt aacaggatta gcagagcgag gtatgtaggc ggtgctacag	3328
agttcttgaa gtgggtggcct aactacggct aactagaag aacagtattt ggtatctgcg	3388
ctctgctgaa gccagttacc ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa	3448
ccaccgctgg tagcgggtgt ttttttgttt gcaagcagca gattacgcgc agaaaaaag	3508
gatctcaaga agatcctttg atcttttota cggggtctga cgctcagtggt aacgaaaact	3568
cacgttaagg gattttggtc atgagattat caaaaaggat ottcacctag atccttttaa	3628
attaataatg aagtttttaa tcaatctaaa gtatatatga gtaaacttgg totgacagtt	3688
accaatgott aatcagtgag gcacotatct cagcgatctg tctatttcgt tcatcatag	3748
ttgctgact c	3759

<210> 15

<211> 42

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 15

ggaggatccg gacggtgagt cactcttggc acggggaatc cg	42
--	----

<210> 16

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 16

ggtgaatatg gctcataaca c	21
-------------------------	----

<210> 17

<211> 23

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 17

ccgccgaaca tggagaacat cgc

23

<210> 18

<211> 33

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 18

cacagatcct ttgttagggt ttaaatgtat acc

33

<210> 19

<211> 29

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 19

ggaggatcct gacctggaag atagtcacc

29

<210> 20

<211> 26

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 20

ggaggatcca tcagcaagca ggtatg

26

<210> 21

<211> 33

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 21

ggagctagcg ggcgtttgac ctccggcgtc ggg

33

<210> 22

<211> 37

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 22

ggagaattca gatctcctct agtaaaacaa tggctgg

37

<210> 23

<211> 25

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 23

ggagctagcc ttotaaccga ggtcg

25

<210> 24

<211> 30

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 24

ggaagatctc cttactccag ctctatgctg

30

<210> 25

<211> 27

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 25

ggcgaattcc ttccgagtga gagacac

27

<210> 26

<211> 43

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 26

ggagtataca tttaaagggc cctaacaaaa caaaaagatg ggg

43

<210> 27

<211> 31

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 27

ggagctagct cgtttacttt gaccaagaac g

31

<210> 28

<211> 36

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 28

ggaagatctc cttatttttg acaccagacc aactgg

36

<210> 29

<211> 29

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 29

ggagtcgacc tgtctgotta cataaacag

29

<210> 30

<211> 26

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 30

cgtaatgctc tgccagtgtt acaacc

26

<210> 31

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 31

gaaagatctc agcaagcagg

20

<210> 32

<211> 47

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 32

ggaggatcct gacctggaag atagtcaggg ttgaggcaag caaaagg

47

<210> 33

<211> 12

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 33

ctagcgggcc ca

12

<210> 34

<211> 12

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 34

gatctgggcc cg

12

<210> 35

<211> 28

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 35

ggagctagca tcatcccagt tgaggagg

28

<210> 36

<211> 28

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 36

ggtagatctc ctcattgtctg ctcgaagc

28

<210> 37

<211> 29

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 37

ccaagctagc gacaaaactc acacatgcc

29

<210> 38

<211> 45

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 38

ggaagatctc gtttaccct gtcatttacc cggagacagg gagag

45

<210> 39

<211> 57

<212> DNA

<213> Artificial sequence

<220>

<223> Oligonucleotide

<400> 39

aagatgtcca gactctgtct ctccgtggcc ctctctgtgc tcctcgggac actcgcc

57

<210> 40

<211> 24

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 40

ggaactagta agatgtccag actc

24

<210> 41

<211> 25

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 41

ggaagctagc ggcgagtgtc ccgag

25

<210> 42

<211> 75

<212> DNA

<213> Artificial sequence

<220>

<223> Oligonucleotide

<400> 42

ggaaagatgg ccagcctctt tgccacattt ctctgtgtgc tcgtgagcct cagcctcgcc
agcgaaagca gcgcc

60

75

<210> 43

<211> 24

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 43

ggaactagtg gaaagatggc cagc

24

<210> 44

<211> 26

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 44

ggaagctagc ggcgctgctt tcgctg

26

<210> 45

<211> 51

<212> DNA

<213> Artificial sequence

<220>

<223> Oligonucleotide

<400> 45

aggtctttgc taatcttggt gctttgcttc ctgccctgg ctgctctggg g

51

<210> 46

<211> 24

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 46

ggaactagta ggtctttgct aatc

24

<210> 47

<211> 25

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 47

ggaagctagc ccccagagca gccag

25

<210> 48

<211> 31

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 48

ggagctagct cgtttacttt gaccaagaac g

31

<210> 49

<211> 25

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 49

ggaagatctc cggtagtggtg tgctg

25

<210> 50

<211> 32

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 50

gcaggatcca gtagacctgg agagaggaca ag

32

<210> 51

<211> 29

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 51

ggaagatcta caaggtgagc tgctgtggc

29

<210> 52

<211> 490

<212> DNA

<213> Pseudo rabies virus

<400> 1

tgccgcgaga ggcggccggg catgcaaatc agaggcgcg	60
gggagacgcc tccgcgcgcc cattggcccg	120
ggcgagccga gatggccgcc ggcggggccg gacatgcaa	180
gtagacgcga gaggaagtag ggagagaaat cccattggcc	240
gtcgaggggc caagatggcg ccctcggggc cggacatgca	300
aagtagacgc gagaggaagt gggcgagaga aatcccattg	360
gccgtcgatg gggcaagatg gccgcgcggg gggccgggca	420
tgcaaatggc cctcgcgagg aagttcctcg cgaatccca	480
ttggccggcg gccgccatct tggccggggc atgcaaagca	490
gacggcagag gaagcgggcy agaaaaatcc cattggcccg	
ccgtcgggga agtcgcgggc gaaaatcggc cattggtccg	
cttacctggg ggcgggctct cctcggggcg cttataagcg	
cgggtctccat cgtagcaatt	

<210> 53

<211> 495

<212> DNA

<213> Rous sarcoma virus

<400> 2

ctgctccctg cttgtgtgtt ggaggctcgt gagtagtgcg	60
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga	120
caattgcatg aagaatctgc ttagggtag gcgttttgcg	180
ctgcttcgcg atgtacgggc cagatatacg cgtatctgag	240
gggactaggg tgtgtttagg cgaaaagcgg ggcttcgggt	300
gtacgcgggt aggagttccc tcaggatata gtagtttcgc	360
taacgatgag ttagcaacat gcottacaag gagagaaaaa	420
gcaccgtgca tgccgattgg tggaagtaag gtggtacgat	480
cgtgccttat taggaaggca acagacaggc ctgacatgga	495
ttggacgaac cactgaatto cgcattgcag agataattgt	
atttaagtgc ctagctcgat acaataaacg ccatt	

ABSTRACT

A nucleic acid construct comprising a chimeric promoter sequence and a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter, wherein the chimeric promoter sequence comprises:

(a) a hCMV immediate early promoter sequence; (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

1/13




Figure 1

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

2/13




Figure 2

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

3/13




Figure 3

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

4/13




Figure 4

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

5/13

Effect of RIA, ENH on SEAP Expression (Two Lines, Three Reps per Line)




Figure 5

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

6/13




Figure 6

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

7/13




Figure 7




Figure 8

9/13




Figure 9

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

10/13




Figure 10

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

11/13




Figure 11

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

12/13




Figure 12

Title: NUCLEIC ACID
CONSTRUCTS
Inventor(s): James T. FULLER
DOCKET NO.: 036481-0141

13/13

Figure 13

ggcgtaatgctctgccagtgttacaaccaattaaccaattctgattagaaaaactcatcgagcatcaaatgaaactgcaatttattcatat
caggattatcaataccatattttgaaaaagccgtttctgtaatgaaggagaaaaactcaccgaggcagttccataggatggcaagatcc
tggtatcggctcgactccgactcgtccaacatcaatacaacctattaatttcccctcgtcaaaaataagggttatcaagtgagaaatca
ccatgagtgacgactgaatccgggtgagaatggcaaaagcttatgcatcttccagacttggtcaacaggccagccattacgctcgtc
atcaaaatcactcgcatacaaccaaacgggtattcattcgtgattgcccgtgagcgagacgaaatagcgatcgctgttaaaggacaa
ttacaaacaggaatcgaatgaacccggcgaggaacactgccagcgcatcaacaatattttcacctgaatcaggatattcttctaata
cctgggaatgctgtttccggggatcgagtggtgagtaaccatgcatacagtgagtagcgataaaatgcttgatggtcggagag
gcataaattccgtcagccagtttagtctgacctctcatctgtaacatcattggcaacgctacctttgccatgttcagaaacactcgg
cgcatcgggcttccatacaatcgatagattgtcgcacctgattgcccacattatcgcgagccattataccataaaatcagcat
ccatgttggaatttaacgcggcctcgagcaagacgttcccgtgtaatatggctcataacacccctgtattactgtttatgtaagcaga
caggctgcacaatattggctattggccattgcatacgttgatctataatataatgtacattatattggctcatgtccaatatgaccgca
tgttgacattgattattgactagtattataatagtaatcaattacggggctcattagttcatagcccatatattggagttccgcttacataacta
cggtaaatggcccgcctggctgaccgccaacgacccccgccattgacgtcaataatgacgtatgttcccatagtaacgccaata
gggacatttcattgacgtcaatgggtggagttttacggtaaaactgccacttgccagtagcatcaagtgtatcatatgccaaagtcggcc
ccctattgacgtcaatgacggtaaatggcccgcctggcattatgccagtagcatgacctacgggactttcctacttggcagtagcatct
acgtattagtcacgtcattaccatgggtgatcggttttggcagtagaccaatggcggtgtagcggttgactcacggggttcca
agtctccacccattgacgtcaatgggagttgtttggcaccaaaatcaacgggactttccaaaatgtcgtataaaccggcccggt
gacgcaaatggcggttaggcgtgtagcgtgggaggtctatataagcagagctcgttagtgaaacgtcagatcgctggagacgc
catccacgctgtttgacctccatagaagacaccgggaccgatccagcctccgcgccgggaacgggtgattggaacgggattc
cccgtgccaagagtgactacccgtccggtatcagcaagcaggtatgactctcagggtggcgcttcccagtcaggaactc
cagggtttagggagcgtgtgggctcttctctacatgtacctttgttgcctcaacctgactatctccaggtcaggatcccagag
tcagggtgctgtatttctcgtgtgtggtcaggtcaggaacagtaaacctgtccgaatattgcctctcacatctcgtcaatctccg
cgaggactggggaccctgtgacgaacatggctagcggggccagatctgggccctaacaacaaaaagatgggggtatttccctaa
acttcattgggttacgaattggaagtgggggacattgccacaagatcatattgtacaaaagatcaaacactgttttagaaaactcctg
taaacaggccattgattgaaagtatgtcaaaggattgtgggtctttgggcttctgtcctcatttacacaatgtggatctcctgcctta
atgcctttgtatgcatgtatacaagctaaacaggctttcactttctcgccaacttacaaggcctttctaagtaaacagtagcatgaacctta
ccccgttgcctggaacggcctgtgtgccaagtgttctgacgcaacccccactggctggggcttggccataggccatcagc
gcatcggtggaacctttgtggtcctctgcccgatccatactgcggaactcctagccgttgttctgcagccggtctggagcaaa
gctcataggaactgacaattctgtcgtcctctcgcggaatatacatcgtttcgtatcgtatgtcttttccctctgcaaaaattatg
gggacatcatgaagcccttgagcatctgacttctggttaataaaggaaatttttattgcaatagtgtgttggaaatttttgtgtct
cactcgggaaggaaattctgcattaatgaatcggccaacgcgcggggagaggcggtttgctattggcgctcttccgcttctcgtc
actgactcgtcgcctcgtcgtcgttgcggcgagcggtatcagctcactcaaaaggcggtataacggtatccacagaatcagg
ggataacgcaggaaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgctgtggtggttttcca
taggtctccgccccctgacgagcatcacaaaaatcagcgtcaagtcagaggtggcgaaacccgacaggactataagatacca
ggcgtttccccctggaagctccctcgtcgtcctcgttccgacctgcccgttaccggatacctgtccgcttctccctcgggaa
gcgtggcgctttctcatagctcacgctgtaggtatctcagttcgggtgtaggtcgttcccaagctgggctgtgtgcacgaacccc
cgttcagcccagccgtcgccttatccggttaactatcgtcttgagtcacacccggttaagacacgacttatgccactggcagcagc
cactggtaacaggattagcagagcgaggtatgtaggcggtgtacagagttctgaagtgggtggcctaactacggctacactagaa
gaacagtatttggatctcgcctcgtcgaagccagttaccttcgaaaaagagttggtagctcttgatccggcacaacaaaccaccgc
tggtagcgggtgttttttgtttgcaagcagcagattacgcgcagaaaaaaggatctcaagaagatcctttgatcttttacggggtct
gacgctcagtggaacgaaaaactcacgttaagggttttggcatgagattatcaaaaaggatctcacctagatccttttaattaaaaa
tgaagtttaaatcaatcaagtatatatgagtaaaacttggtcgtacagttaccaatgcttaatcagtgaggcacctatctcagcgatct
gtctatttcgtcatccatagttgcctgactc